

**Responses shown by bird
communities to teak plantations in
Sagar Forest Division, Karnataka**

*Dissertation Submitted to
University of Saurashtra, Rajkot*

*In partial fulfilment of
Master's Degree in Wildlife Science*

By

SAHAS S. BARVE

Under the Supervision of
Dr. Dhananjai Mohan and Mr. Qamar Qureshi



**भारतीय वन्यजीव संस्थान
Wildlife Institute of India**

June 2009



Certificate

This is to certify that Mr. Sahas S. Barve has carried out an original piece of research in partial fulfilment of Master's Degree in Wildlife Science of the Saurashtra University, Rajkot. The topic of his dissertation is "Responses shown by bird communities to teak plantations in Sagar Forest Division, Karnataka". The investigations were carried out under our supervision from November 2008 to June 2009. We hereby certify that this work has not been submitted for any degree to any university.

Dr. Dhananjai Mohan
Supervisor

Sh. Qamar Qureshi
Co-Supervisor

Date: 3-08-09

Place: Dehradun

Contents

<i>List of Tables</i>	<i>i</i>
<i>List of Figure</i>	<i>ii</i>
<i>Acknowledgements</i>	<i>iv</i>
<i>Summary</i>	<i>vii</i>
1. Introduction	1
1.1 Literature review	2
1.1.1. Choice of plantation species	2
1.1.2 Age of plantation/stand	3
1.1.3 Landscape matrix	3
1.1.4 Proximity to natural forest	4
1.1.5 Plantations as bird habitats	4
1.1.6 Teak plantations- a brief history and role in biodiversity conservation	6
1.2 Objectives	8
2. Study area and Design	9
2.1 Location	9
2.2 Climate and Rainfall	10
2.3 History of Forestry	12
2.5 Vegetation	12
2.6 Biodiversity	13
2.7 Human disturbance	13
2.8 History of teak plantation forestry in the division	14
2.9 Study design	15
2.10 Selection of sites for sampling	16
3. Methods	17
3.1 Bird Sampling	17
3.2. Vegetation Sampling	18
3.3 Analysis	20
4. Results	25
4.1 Density of Birds	25
4.1.1 Comparisons of Overall densities	26
4.1.2 Comparisons between strata within seasons	26
4.1.3 Comparisons between strata across seasons	27

4.1.4 Comparison of abundances according to habitat guilds across seasons	29
4.2 Diversity of Birds	30
4.2.1 Estimated species richness winter	30
4.2.2 Estimated species richness summer	32
4.2.3 Diversity estimation for transects	34
4.3 Community Similarity	35
4.3.1 Morisita Horn Index comparisons	35
4.3.2 Sorenson Index comparisons	36
4.4 Vegetation profile of the three strata	36
4.4.1 Structural attributes of the three strata	36
4.4.2 Floristic attributes of the three strata	41
4.4.2.1 Estimated richness of trees in the strata	41
4.4.2.2 Diversity of trees and shrubs in the strata	41
4.5 Habitat Correlates explaining bird diversity	43
4.5.1 Principal Component Analysis	43
4.5.2 Pearson's Correlations for bird responses to habitat attributes winter	43
5 Discussion	46
5.1 Differences in habitat characteristics	46
5.2 Density of birds	47
5.3 Diversity of birds	51
5.4 Community structure	53
5.5 Habitat attributes explaining bird density and diversity responses	54
5.5.1 Patterns common to both seasons	55
5.5.2 Seasonal patterns	55
5.6 Conclusion	57
Literature cited	59

LIST OF TABLES

Table 2.1 List of mature plantations selected for bird sampling	16
Table 2.2 List of Young plantations selected for bird sampling	16
Table 4.1- Overall (strata level) density estimates (Birds/ha), the values in parentheses denote % coefficient of variance.	22
Table 4.2- Guild level density estimates (Birds/ha) in winter - the values in parentheses denote % Coefficient of Variance	22
Table 4.3- Guild level density estimates (Birds/ha) in summer - the values in parentheses denote % Coefficient of Variance	22
Table 4.4 - Estimated species richness (winter) using JACK 1 estimator	32
Table 4.5- Estimated species richness (summer) using JACK 1 estimator	34
Table 4.6- Shannon's Index values for each transect in the two seasons	34
Table 4.7- Community similarity using Morisita Horn Index winter. Values in parantheses denote number of species shared between the two strata	35
Table 4.8- Community similarity using Morisita Horn Index summer. Values in parantheses denote number of species shared between the two strata.	35
Table 4.9- Shannon's index for tree and shrub diversity for transects	41
Table 4.10 Contribution of each variable to the extracted factors	43

LIST OF FIGURES

Figure 2.1-	Map of Study Area- The highlighted ranges is the intensive study area	10
Figure 2.2-	Study Design	14
Figure 3.1	Box plots showing median distance for detections of the 4 size classes analyzed in winter (a) Natural forests (b) Mature plantations (c) Young plantations	20
Figure 3.2	Box plots showing median distance for detections of the 4 size classes analyzed in summer (a) Natural forests (b) Mature plantations (c) Young plantations	20
Figure 3.3-	Hierarchical classification of data used in analysis.	21
Figures 4.1-	Graphs showing proportion of each guild in total density in Natural forests stratum across seasons.	27
Figure 4.2	Graphs showing proportion of each guild in total density in mature plantations stratum across seasons.	27
Figure 4.3-	Graphs showing proportion of each guild in total density in young plantations stratum across seasons.	27
Figures 4.4-	Graphs showing proportion of detection of each habitat guild in 15m belt in transects in natural forests.	28
Figures 4.5-	Graphs showing proportion of detection of each habitat guild in 15m belt in transects in mature plantations.	28
Figures 4.6-	Graphs showing proportion of detection of each habitat guild in 15m belt in transects in young plantations.	28
Figure 4.7-	Overall estimated species richness winter	29
Figure 4.8-	Estimated species richness Insectivorous guild winter	30
Figure 4.9-	Estimated species richness for phytophagous guild winter	30
Figure 4.10-	Overall estimated species richness summer	30
Figure 4.11-	Estimated species richness insectivorous guild summer	32
Figure 4.12-	Estimated species richness for phytophagous guild summer	32
Figure 4.13-	Graph of Sorenson index of similarity between strata in winter and summer.	35
Figure 4.14-	Distribution of foliage in a vertical spread of vegetation	36

Figure 4.15- Variability in crown height of trees in each transect.	36
Figure 4.16- Difference in vertical spread of foliage in the three strata.	37
Figure 4.17- Distribution of Girth at Breast Height in trees in the three strata.	37
Figure 4.18- Variability in GBH in trees of the three strata.	38
Figure 4.19- Tree density/m ² in the three strata.	38
Figure 4.20- No. of parasites/ vegetation sampling plot	39
Figure 4.21- Seasonal difference in canopy cover in the three strata	39
Figure 4.22- Estimated tree species richness using JACK 1 estimator	40
Figure 5.1- Increase in abundance of birds common to both seasons in natural forests	46
Figure 5.2 - Increase in abundance of birds common to both seasons in mature plantations	47
Figure 5.3 - Increase in abundance of birds common to both seasons in young plantations	47
Figure 5.4- Rank abundance of phytophagous birds in young plantations in winter.	49
Figure 5.5- Rank abundance of phytophagous birds in young plantations in summer.	50

Acknowledgments

I would like to thank the Director, Dean of Faculty, Wildlife Sciences and all the faculty members of the Wildlife Institute of India for their invaluable guidance and support throughout the course and dissertation.

I would like to thank Sh.Dhananjai Mohan and Sh. Qamar Qureshi, my supervisors who were the constant guiding lights through every stage of my dissertation. Dr. Dhananjai Mohan for some very useful inputs in field and with analysis and Sh Qamar Qureshi for his invaluable guidance in analysis and for making me see my data in new light.

Special thanks to the Karnataka Forest Department, Sh. Sanjai Mohan, Conservator of Forest Shimoga Circle, Mr. Mahesh Shirur, Divisional Forest Officer, Sagar Forest Division, and all other staff for their ample cooperation.

I thank Dr. T.R. Shankar Raman and Dr. Rajah Jayapal for helping me conceptualize my dissertation topic and going through my proposal and giving some very valuable suggestions. Abishek, Mousumi, and Upamanyu Hore (Dada) for some practical tips in structuring my proposal and Hari Shridhar, Raman and Navendu Page for helping me find some important literature.

I would like to thank Dr. Gururao Bapat and Mrs. Madhuri Bapat for their warm hospitality, great food and enriching conversations, I thank them for keeping their huge collection of books open to me at all times. I cannot thank them enough for making me SO comfortable in a place and at a time when I was doing so many things in a place with a huge language barrier. I thank Sujendra Kumar K.C.(Suji) for being a great friend and a great help with field work. I have learnt many things from his unique outlook towards life. I also want to thank Mani anna, Navin anna, Satish anna and Dnyana madam for accepting me and helping me throughout my time in Sagar. I thank Prakash anna and family from Bacchhodi for their hospitality and help. Saiju

anna for taking me on night trails for civets and flying squirrels, Bindoo akka for delicious food and Masterji for translating all the kannada to hindi. Saiju anna and Suji were also immense help in finding the Ceylon frogmouths in Kogar. I want to thank Mr. Hari Divekar and Mr. Rama Prasad for logistic help in Kogar and Shettyhalli. I also want to thank Navendu Page, for identifying the diverse flora of my study area. Varun Torsekar and Ishan Agarwal and Navendu again for a splendid frogging trip. A huge thanks to Raviraj for visiting me in field. I thank the shamas for filling every morning in field with their fluty songs and the collared scops owl for making my nights magical .

I want to thank Abhi, Moss, Rekha, Sartaj, Chandan, Nilanjana Madam, Mr. Monoj Agarwal, Mr. Manoj Nair and Pranav Chanchani for help in analysis and thesis writing. I thank Deep Contractor and Jennie for going through the many versions of my thesis and giving their useful comments. Here I would especially like to thank Sartaj and Rekha for some mind numbing but greatly enlightening conversations on fundamentals of statistics.

Back in the institute I want to thank Murali for introducing me to and being a partner in crime for everything bad, .and for the (most useless and unwanted) trivia. Asif for being a fun room mate in all the field visits and those lovely dinners of kebabs and momos. Rekha for being a hard core Mumbaite and a firm support throughout the course. Sartaj, for the many “haiku” moments which were followed by insane bouts of laughter. Dutta for his generosity and being an amazing friend and fellow birder. Sumithra for having the exact opposite taste as mine in everything in the world and still being able to have so much fun together. Upamanyu Hore (dada), Merwyn Fernandis, Udayan, Bibek (Khamba), Bopanna, Kausik and Sutirtho for their incessant support and for the intense, sometimes toxic debates over lunch, tea and other things about conservation amongst other things.

I would like to thank Dr. G.S. Rawat for his invaluable field inputs, his never tiring nature for botanically challenged people like me, the thrilling badminton games and above all for being an excellent teacher . Dr. Goyal, Dr. Uniyal, Sh. Qamar Qureshi, Dr. S.A. Hussain, Dr. Sathyakumar, and Dr.Adhikari for some memorable time in field. Dr. Vasudevan, Dr. Jhala Dr. Goyal for some highly invigorating conceptual classes. Dr.Shankar, Dr. Malik and Dr. Nigam, Dr. Shivkumar, Sh. Dhananjai Mohan Sh. A. Udayan, Dr. B.K. Mishra and Dr. R. Badola for making some seemingly boring subjects, very interesting.

I would like thank Ashutosh sir, Dhamanda sir and all the academic cell staff. Shambhuji and Gulshan for their prompt help and ever smiling face for making our days in WII extremely comfortable. Mr. Rajesh Thapa, Mr. Verrappan and all the staff of the computer department for their support.

Dr. Hussain and Pathakji for hostel facilities and an extremely accommodating nature. I would like to thank Rehan Aunty for her warmth and being an excellent guardian, Dr. Ghuman (Sartaj's mom) for the absolutely divine apple pies amongst other things. Khamba and Aunty in Lhasa restaurant for breaking the routine of repetitive hostel food with some gastronomic delights. I would like to thank Dinesh and Rakesh from the Old Hostel Mess for being ever smiling at breakfast.

I want to thank my batch mates in the XIth MSc course- Sartaj, Rekha, Murali, Asif, Datta Sumithra, Chandan and Sabita for making these two years, the best two years of my life. They have been a constant source of inspiration, laughter and pure delight. Finally I thank my parents, my brother, Shamoli, and all my friends back home Milind, Rahul, Raviraj, Suruchi, Isha and Aditya amongst others.

Summary

This study was aimed at discerning the responses shown by bird communities to teak plantations of various age classes in Sagar Forest Division, Karnataka within Latitude $13^{\circ} 36'$ and $14^{\circ} 38'$ North 74° Longitude $38'$ and $75^{\circ} 32'$ East Latitude. Birds were sampled using line transects in three habitat strata, natural plantations, mature teak plantations and young teak plantations. Density, diversity and community similarity were the chief ecological responses that were quantified to bring out differences in the usage of forest plantations by native bird fauna. Habitat (vegetation) correlates, both structural and floristic were quantified to establish the reasons for differences. The sampling period was from December 2008 to April 2009. This period was divided into two seasons, winter and summer. A total of 9 transects were laid, marked and sampled at least 4 times per season.

Birds were divided into three feeding guilds and three habitat guilds for analysis. The overall density of birds across the three strata was found to be higher in summer than in winter. However a guild level analysis showed that in winter while insectivorous birds were most abundant in natural forest with decreasing density towards young plantations ($F_{(6,241)}$, $p < 0.05$). There was no difference in their density in summer. Phytophagous birds showed the exact reverse trend. Density of this guild was higher in young plantations than natural forests and mature plantations in winter and summer ($F_{(5,689)}$, $p < 0.05$), ($F_{(14,302)}$, $p < 0.01$) respectively. Overall diversity of birds showed a trend of decreasing species richness from natural forests to young plantations in both seasons. However this difference was almost non existent in summer at the guild level. The abundance of birds belonging to evergreen and moist deciduous forests shows a consistent rise in abundance from winter to summer in all the three strata. There is also evidence that there is ingression of individuals of species that were

common to both seasons possibly for nesting. The overall community similarity is quite high between the three strata and it increases from winter to summer. Bird diversity was found to be affected by the vertical spread of vegetation and tree height heterogeneity ($R=0.871$, $p<0.01$). It was also correlated to canopy cover ($R=0.807$ $p<0.05$) and continuity ($R= 0.737$ $p< 0.01$). Insectivorous guild was affected by correlates of forest maturity ($R= 0.805$ $p< 0.01$). The high density of phytophagous birds was found to be correlated to the abundance of parasitic epiphytes which were most numerous in young plantations ($R= 0.766$, $p< 0.05$ in winter and $R= 0.796$, $p < 0.01$ in summer). Although resident bird composition showed association with tree species composition in winter ($R= 0.49$, $p< 0.001$), no such association was observed in summer.

1. Introduction

As rates of deforestation continue to rise in many parts of the world, the international conservation community is faced with the challenge of finding approaches which can reduce deforestation and provide rural livelihoods in addition to conserving biodiversity. However, much of modern day conservation is motivated by a desire to conserve 'pristine nature' in protected areas (Bhagwat et al 2008). The burgeoning human population has already removed somewhere between 8 and 12 million square kilometres (between 35% and 50%) of the original closed canopy tropical forests around the world (Wright and Miller-Landau 2006). The loss of natural forest is most profound in the tropics where annually 0.8% of the remaining forests are cleared and converted into cultivated land. Growing human populations and accelerating demand for forest resources such as wood resulted in increasing areas of degraded forests and forest plantations. New forests are regenerating on former agricultural land, and forest plantations are being established for commercial and restoration purposes (Nagendra 2007, Chazdon 2008). During the past thirty years an evident decline in natural forest resources in a number of countries and the difficulties in accessing increasingly remote areas of natural forest available for wood supply, has resulted in intensification in the focus on plantation forests. Plantations provide a potential means for alleviating potential future wood shortages and providing continuity of supply for existing industrial enterprises or household fuel wood needs (FAO 2002). Apart from fuel and timber biomass, plantation forests are also known to provide services like carbon sequestration and soil conservation.

Role in biodiversity conservation: Although networks of protected areas in the tropics provide habitat for many rare and endemic species that prefer old-growth forest, these networks alone fail to protect all tropical biodiversity due to their small

land area (Montagnini et al 2005). Plantation systems can buffer existing reserves and provide corridors for persistence and movement of species across landscapes (Daily et al 2001). Such systems are useful with pristine forests for combating species loss as a result of tropical forest fragmentation.

1.1 Review of Literature

Birds in plantations: Birds are one of the most well studied groups of organisms. With their varying levels habitat specificity, birds make excellent indicators of ecosystem health and habitat quality. Hence, birds can be used to assess the quality of habitats the growing plantations offer and use them as a surrogate for biodiversity. Biodiversity conservation is thought to be an inherent service provided by the plantations of the world. But it is important to understand what aspect of biodiversity is being conserved. An increasing number of studies suggest that the differences in bird communities in the plantations versus natural forests are quite stark. Generally not more than 50-60% of the original bird fauna is supported (Bhagwat et al 2008). Petit et al (1999) reported a mere 43%-54% forest bird species present in rustic coffee plantations. Similar results have been shown from Costa Rica (Daily et al 2001), India (Raman 2001), Kenya (Sekercioglu 2002) and many other studies. Research indicates that in the plantations forest specialist birds are most commonly replaced by more generalist, open habitat species that may not be of significant conservation importance (Kwok and Corlett 2000, Daily et al 2001, Rotenburg 2007).

1.1.1 Choice of species for plantations: In a study from Kenya Farwig et al (2008) illustrated that plantations of indigenous trees support more bird diversity than those of exotic ones. Plantations of a mixture of native species more closely resembled natural forests with respect to bird species composition than monocultures of a single species. The bird conservation significance of using native tree species has also been

reported from studies on rustic (native) shade tree coffee plantations in the Americas (Greenberg et al 1998, Moguel and Toledo 1999, Jende et al 2005, Tezeda-cruz and Sutherland 2006,) and India (Anand et al 2008). The choice of species for plantation is known to be extremely important for certain guilds of birds like frugivores and cavity nesting birds (Clout and Gaze 1984).

1.1.2 Age of stand/plantation: The age of the plantation is also important in determining the usage by local bird communities. This trend has been seen with *Tectona grandis* plantations in India, in which the older plantations always seem to have a greater diversity of birds and a more stable community structure than the younger monocultures (Daniels 1989, Mehta 1998, Trivedi 2006). Because plantations are generally man-made habitats, it takes many years for a stable bird community to establish. Hence it is recognized that plantations of tree species with long rotation cycles provide time for the system to stabilize and provide the habitat complexity needed for higher bird diversity.

1.1.3 Landscape matrix: There is an increasing body of literature on the importance of the landscape matrix in determining bird diversity of plantations. A landscape has been defined as an area with an interaction of its elements (e.g. ecosystems) relevant to some phenomenon under consideration (Mazerolle and Villard 1999). Many bird species react to small scale (patch level) and landscape level changes in habitats including the ones which extend beyond their home ranges (Taylor and Krawchuk 2005). Plantations alone might not support a large diversity but where they are located in close proximity to natural forest, they can give good cover for many species and hence can act as good buffers for these habitats especially for dispersing individuals.

1.1.4 Proximity or connectivity to natural forest: Raman (2006) reports that the species diversity and richness of some native tree plantations located adjacent to forest fragments were comparable to wet evergreen forest habitats. But as the distance of the plantation increased away from the natural forest vegetation, the number of true forest species declined. Lawes and Wethered (2005) in their study of the plantation forest matrix in Afromontane ecosystem describe the importance of canopy connectivity. The importance of canopy connectivity has also been realized in forest fragments- agro-ecosystem network in Panama (Petit et al 1999), and the Israeli scrub desert (Shochat et al 2001) also. Hence it is evident that not only the existence but also the pattern of juxtaposition of forest plantations and natural forest areas is important in conserving forest bird communities.

1.1.5 Plantations as bird habitats: The composition of avian assemblages and how they are influenced by habitat features has been one of the most pervasive themes of investigation in community ecology (Jayapal et al 2009). Habitat structure (physiognomy) and floristics (composition) have been recognized as the key components affecting land bird assemblages in general. It is important to assess plantations using these two attributes for their value as bird habitats.

Physiognomy: Birds are known to respond to two broad aspects of habitat structure, vertical heterogeneity and horizontal or spatial heterogeneity. While Foliage Height Diversity and Tree Height Diversity are important attributes of vertical heterogeneity, canopy cover and spread, tree density, number of snags, basal area, shrub density and volume are determinants of spatial heterogeneity (MacArthur and MacArthur 1961). Plantations because they are planted in a uniform fashion and very seldom regenerated naturally offer very little structural heterogeneity. Tree density is generally known to be higher in plantations than in natural forests (Saha 2003,

Daniels 1992). Sekercioglu (2002) reported that there was definite distinctness between the bird communities found in unlogged or little logged forests and heavily logged forests. This pattern, he explained due to the loss of horizontal heterogeneity. Hansen et al (1995) and Wilson (1974) reported the relationship between the density of snags and dead poles and their effect on the abundance of cavity nesting birds. The loss of these nesting sites seriously affected this guild and there was a marked recovery when snags were retained. Marsden et al (2001) reported that Eucalyptus plantations lacked substantial bird fauna as there was little or no under storey vegetation in these plantations.

Floristics (composition): This aspect of habitat heterogeneity is known to play a significant role when comparing sites in the same overall habitat type (Lee and Rottenburry 2005). Bell (1979) reported very little native plant and bird diversity from teak plantations in Papua New Guinea. Arnold (2003) showed that Australian agro-forestry areas showed that the difference in under storey diversity translated into a difference in both diversity and abundance of bird species.

1.1.6 Teak plantations- Brief history and studies on their role in biodiversity conservation.

Teak (*Tectona grandis*) is a hardwood species indigenous to India. Teak has been grown in other countries including Costa Rica and Papua New Guinea, Central and South America and Africa where it is an exotic making it the most widely planted high quality timber species covering about 2.3 million ha (Enters 2000, F.A.O., 2002). Teak plantations exist in many parts of India and often replace natural evergreen forests and also moist and dry deciduous forests. Today it has the second biggest share in the national growing stock contributing 6.31% of the stems and 4.33% of the volume of trees in forests (SFR 2005).

Bell (2000) reported that in Papua New Guinea, Teak had little lateral branching, few epiphytes or climbers and little undergrowth. There were far fewer species of birds and mammals than in the rain forest, owing mainly to the loss of small insectivorous passerines, nectarivores and ground feeders. Teak plantations of different age classes especially young plantations if intermixed with natural forest provide good habitat to large mammals in the Western Ghats of India. (Nair and Jayson 1989, Harikrishnan et al 2007).

Few studies have been done on birds found in teak plantations in India. Prachi Mehta (1998) in her study of birds in different age classes and silvicultural systems of teak in Bori Wildlife Sanctuary, found that the diversity differed little after a teak plantation had reached 20 years. The density of birds however showed considerable variation in clear felled areas and teak plantations. Daniels (1989) worked on birds of Uttara Kannada and demonstrated that the teak plantations were very homogenous habitats and supported a considerable proportion of the local avian species fauna. Trivedi (2006) reported that species richness of birds increased from young monocultures of

teak to moist deciduous forests in the Dangs. Hence teak plantations usually follow the same trends in diversity as most other plantation crops.

The present study compares density and diversity of birds in teak plantations of two age classes with the bird community in the surrounding natural forests. It examines bird abundance and diversity at a guild level across the three strata. A vegetation profile of the three strata is carried out to highlight differences in structural and floristic attributes and describing the vegetation of the study area. Finally the study investigated habitat attributes that explain bird community responses to teak plantations.

1.2 Objectives

- 1) To estimate density and diversity of birds in teak plantations in Sagar Forest Division, Karnataka and the surrounding natural forests.
- 2) To evaluate the extent of similarity in bird community structure and composition of teak plantations and surrounding natural forests.
- 3) To find habitat correlates which explain the differences in density and diversity of birds in these strata.

These objectives are used to answer the following research questions-

- 1) What are differences in overall density of birds in natural forests, mature teak plantations and young teak plantations in winter and summer?
- 2) What are differences in density of birds at the guild level across the three strata in winter and summer?
- 3) What are the differences in species richness as a measure of diversity among the three strata across two seasons?

- 4) What are the bird attributes that explain the seasonal difference in density and diversity?
- 5) What is the level of similarity/dissimilarity in the bird community of the three strata?
- 6) What are the differences in the structural and floristic attributes of the vegetation of the three strata?
- 7) What are the habitat variables that explain the differences in the bird density and diversity in the three strata in the two seasons?

2. Study Area

The study was carried out in the Sagar Forest Division situated in the Shimoga district in the state of Karnataka. The division is named after the Sagar town that houses the headquarters of the division. The division is divided further into ten ranges (Murthy, S.F.D. Working plan 2001). The current study was carried out in the ranges of Sagar, Anandpuram and Soraba.

2.1 Location

Sagar Forest Division is situated within 13° 36' and 14° 38' North 74° 38' and 75° 32' East. The total area of the division is 14,782 km². The area is very hilly and undulating. The highest point in the division is Kodchadri with an elevation of 1341m with the average elevation is 670 m M.S.L. The significant rivers flowing through the division are the Sharavathy, Kumudwathi, Varada, Varahi and Chakra. Water availability generally does not fluctuate seasonally (Murthy, S.F.D. Working plan 2001).

2.2 Climate and Rainfall

The climate of the area is moderate with temperatures varying between 13°C and 37°C. The Southwest monsoon brings most of the rain, commencing usually in the early part of June with very short interruptions. Heavy and continuous rainfall is received until September with an average annual rainfall of 2400mm. Pre- monsoon showers arrive in May. November, December and January are the coolest months. Summers are severe with March-May being the hottest months of the year.

The world famous Jog Falls on the course of the Sharavathy River are present in the division. Two Protected Areas, Gudavi and Sharavathy Wildlife Sanctuaries also fall within the limits of the division (Murthy, S.F.D. Working plan 2001).






Figure 2.1 Maps of Study Area- The highlighted ranges is the intensive study area and location of transects within those ranges.



2.4 History of Forestry

This area has a rich history of forestry dating back to the late nineteenth century. Although the division came into existence for the first time in 1916, but it has maintained a permanent division status only since 1932. The oldest reserved forests of the area were given protection in the year 1888. The area has been a source of massive extractions of many commercially important timber species such as *Tectona grandis*, *Pterocarpous marsupium*, *Terminalia alata*, *Terminalia paniculata*, *Lagerstromia lanceolata*, *Haldina cordifolia* and *Xylia xylocarpa*. Minor Forest Produce of the area includes bamboo and sandal wood.

2.5 Biodiversity

Vegetation

The forests of the area can be divided into four main categories (Champion and Seth 1968). These are as follows-

- 1) Southern Tropical Wet Evergreen Forest
- 2) Southern Tropical Semi-Evergreen Forest
- 3) Southern Tropical Moist Deciduous Forest
- 4) Southern Tropical Dry Deciduous Forest

The wet-evergreen forests are limited to the Hosnagara, Kargal and Nagar ranges. Due to deterioration by biotic pressures, the forests of the area show a tendency to change from wet-evergreen forests to semi-evergreen forests and from semi-evergreen to moist deciduous forests.

The present study was carried out in the moist-deciduous forests of the Sagar, Anandpuram and Soraba ranges. The semi-evergreen and moist-deciduous forests of the area are not very dense. Although an intimate mixture of species is observed, only

a few species make up the canopy. The chief feature of these forests of the area is that they do not shed leaves until well into the dry season, which is late February or March. Interestingly, they come into new leaf long before the monsoon starts. The important species of this zone include *Tectona grandis*, *Pterocarpous marsupium*, *Terminalia alata*, *Terminalia paniculata*, *Lagerstromia lanceolata*, *Haldina cordifolia*, *Xylia xylocarpa*, *Anogeissus latifolia*, *Dalbergia latifolia*, *Bombax ceiba*, *Mangifera indica* (Murthy, S.F.D Working Plan 2001)

Fauna-

Mammals- The reserved forests do not have a high density of large mammals, chital (*Axis axis*), wild pig (*Sus scrofa*) and sambar (*Cervus unicolor*) are occasionally encountered. Bonnet macaques (*Macaca radiata*), black napped hare (*Lepus negricollis*) hanuman langur (*Seminopithecus entellus*), jackal (*Canis aureus*) and Indian giant squirrel (*Ratufa indica*) are more common(Pers. Observ.)

Birds- The division is rich in bird life. Although the study was aimed at forest birds and was limited to communities in moist deciduous forests where a total of 119 species were observed, visits to other localities and habitats in the division yielded a total of 247 species.

2.7 Anthropogenic Pressures on Wildlife Habitats: The area is not thickly populated and hence the human impact on the forests is not very high (Murthy A., S.F.D. Working Plan 2001). Nonetheless a large amount of biomass is extracted in the form of firewood and leaf litter. Firewood is used in most homes and hotels particularly in Sagar town, hence generating a large demand on the forest. Leaf litter collection is done in the summer season using sweeping techniques leaving the ground almost bare. The collected litter is mixed with cattle dung as an organic

fertilizer in the areca nut (*Areca catechu*) plantations that occupy most of the low-lying areas in the region. Hunting is not a significant problem in the area, although the sparse density of mammals hints at a probable past of hunting in the area.

2.8 History of Teak Plantation Forestry in the Division:

Sagar division has 7815.6 ha of teak plantations. A teak conversion working circle manages the teak plantations within the division. The main objective of the circle is to exploit the marketable timber and conversion of existing growing stock into valuable teak plantations. Ten felling series were formed, with a rotation period of eighty years and an exploitable girth of 4.5 feet. Though some thinnings were carried out in the past they depended mainly on the availability of funds. Due to irregular and unsystematic workings in the past, Laurey's thinning formula for teak plantations has been adopted. The main objectives of management are-

- 1) To convert existing teak plantations into valuable growing stock by following proper thinning schedule.
- 2) To enrich plantations with concentrated artificial regeneration of teak if they are damaged due to biotic pressures.
- 3) To meet the teak pole requirement of the local and distant stakeholder.
- 4) To convert plantations into mixed nature if teak has failed on account of some locality factors.
- 5) To remove Eucalyptus coppice and convert areas into pure teak stands.

According to the existing silvicultural practices there are six thinnings in one rotation cycle and the number of stems retained per hectare changes from 1250 per hectare to 198 per hectare after the sixth thinning in the 48th year. (Murthy, Working Plan S.F.D. 2001).

2.9 Study Design:

There were three strata namely- Natural forests, Mature plantations and Young plantations. Each stratum had three transects in each.




Figure 2.2 Study Design

2.10 Plantations chosen for the study: The plantations for the study were selected after a weeklong survey of the area. They were chosen based on whether transects could be laid in the plantations while maintaining a buffer of at least 150 m from each side and that fell in the desired age class category. Care was also taken that the plantations were primarily surrounded by moist deciduous forests to control for effects of surrounding habitats on the bird community within the plantations.

Table 1.1- List of mature plantations selected for bird sampling

Name of Plantation	Year of establishment	Area (ha)
Ullur	1965	41
Konehosur	1962	140
Malali	1962	50

Table 1.2- List of Young plantations selected for bird sampling

Name of Plantation	Year of establishment	Area (ha)
Teliganmane	1989	40
Bilisiri	1987	50
Konehosur	1987	65

Hence the average age of mature plantations is 47 years and that of young plantations is 23 years.

Natural Forest:

Three natural forest patches were also selected for sampling. These patches were chosen on the basis of their quality as bird habitats and distance to the selected teak plantations (≤ 3 km). The quality of these patches was assessed by carrying out reconnaissance surveys in the early mornings and making multiple bird species richness lists. The patches were also chosen in a fashion so as to control for distance of teak plantation to closest natural forest.

3 Methodology

3.1 Bird Sampling-

Distance sampling (line transects)

Birds were sampled using variable width line transects (Rosenstock et al 2002, Thomas et al 2002). Line transects were chosen as they give maximum detections/unit time in field and hence effectively cover more area in the same time as compared to point counts (Verner 1985). The basic assumption of line transects is that all objects on the line are detected and the probability of detection goes on decreasing as distance from the line increases. Hence a detection function is fitted on the observations so as to arrive at the required abundance estimates.

Three transects each were laid out in three strata mature plantations, young plantations, and natural forests. All transects were between 500m-700m in length. The transects were laid in a way so that there was a buffer of at least 150m from all sides to negate 'edge effects'. All plantations that were chosen for sampling all had semi evergreen and moist deciduous forest on at least three sides so as to control for 'neighbourhood effect' (Wethered and Laws 2005)

All transects were walked at a steady pace (approximately 1.25 km/hour) between 6:30 a.m. and 9:30 a.m. At each detection of a bird species the following variables were recorded- 1) Species 2) Group size 3) Angle between the observer and the bird using a standard hand held compass 4) Distance between the observer and the bird using a laser rangefinder 5) The height class at which the bird was seen. The birds were recorded as being in one of 4 height classes. If they occurred in the first 0-5 m above ground then they were recorded in height class 'a'. If between 5m-10m then in height class 'b'. If between 10m-15m then in height class 'c' and if between 15m and above then in height class 'd'. These height categories were selected after

reconnaissance walks in the plantations and developing a visual perception of the heights at which vegetation was stratified. Similarly rough height classes were chosen at heights where the birds were observed. Only birds actually perching in the habitat were recorded. Birds were also detected and recorded using aural cues. On detection by the call of the bird, the angle and distance to the tree or bush the call originated from, was recorded.

Sampling was divided into two seasons- winter and summer. Each transect was walked a minimum of 4 times (pseudo-replicates) in winter and a minimum of 5 times in summer, with a total of 49 walks in winter and 57 walks in summer combined in all strata.

3.2 Vegetation Sampling

The following habitat attributes were quantified as covariates to explain bird community responses to teak plantations. The variables were classified into two main categories-

- 1) Structural Covariates: These quantify the structural attributes of the habitat.
- 2) Floristic Covariate: These quantify the floristic attributes.

Structural Attributes

1. **Tree density:** Number of live trees was counted. A tree is defined as any woody plant > 2m in height (Jayapal 2007).
2. **Girth at breast height (G.B.H.):** G.B.H of all trees in the plot was recorded.
3. **Tree height diversity:** The lowest and the apical height of branches of the trees was recorded and the Shannon Weiner Index was derived $-\sum [P_i * \ln(P_i)]$ where P_i = proportion of trees in the i^{th} height class (Jayapal et al 2007, Mohan 2007)

4. **Canopy cover and spread:** Percentage canopy cover as measured by a spherical densiometer. Mean of four directional readings were taken from the centre of the plot.
5. **Shrub abundance:** Number of shrubs was counted in each plot. For the purpose of this study, a shrub is defined as any woody plant < 2m in height.
6. **Shrub volume:** Mean shrub volume as the product of height X breadth X width was recorded for each shrub wherever possible.
7. **Ground cover:** Ocular estimation of % grass cover, % litter cover and % bare soil in each plot was recorded.
8. **Parasites:** Number of plant parasites present on trees in the 10m circular plots was recorded.
9. **Bamboo clumps:** A total count of bamboo clumps was done in a belt of 60m on both sides of the transect.

Floristic attributes

- 1) **Tree species richness:** Number of tree species encountered in each plot was recorded.
- 2) **Tree species diversity:** This was calculated by using the Shannon's Index as $-\sum [P_i * \ln(P_i)]$ where P_i = proportion of trees belonging to species 'i'.
- 3) **Shrub species richness:** Number of species of shrubs encountered in each plot was recorded.
- 4) **Shrub species diversity:** This was calculated by using the Shannon's Index as $-\sum [P_i * \ln(P_i)]$ where P_i = proportion of shrubs belonging to species 'i'.

3.3 Analysis

Density of birds

We checked for a size bias in the distance sampling of small sized birds. This was done by dividing the birds into 4 size classes. Birds up to 15 cm in size were classified into size class '1'. Species that measure between 15 cm -20 cm in length were classified as group '2'. Species up to 25 cm in length were classified as group '3'. All bird species larger than 25 cm were placed in group '4' (Ali and Ripley 1973).

Winter




Fig 3.1 a Winter Natural




Fig 3.1 b Winter Mature




Fig 3.1 c - Winter Young

Summer




Fig 3.2 a Summer Natural




Fig 3.2 b Summer Mature




Fig 3.2 c- Summer Young

Figure 3.1 and 3.2 Box plots showing median distance for detections of the 4 size classes analyzed in winter and summer respectively (a) Natural forests (b) Mature plantations and (c) Young plantations

When no size based bias was observed then all the bird detections were used to carry out density estimations.

Density of birds was estimated using the programme DISTANCE 5 (Thomas et al. 2006). The programme allows for global estimates (of the entire data set) of density and also allows stratification according to sampling design and also ‘post stratification’ for known subsets of data. In DISTANCE models datasets for clustered observations are modelled separately than those for single object observations (Rosenstock et al. 2002). Transect datasets were analyzed in the following manner-

Season

Winter	Summer
--------	--------

Habitat Strata

Natural	Mature	Young
---------	--------	-------

Bird Feeding Guild

Insectivorous Guild	Phytophagous Guild	Others Guild
------------------------	-----------------------	-----------------

Figure 3.3 Hierarchical classification of data used in analysis.

Birds were assigned to three basic feeding guilds- 1) Insectivorous 2) Phytophagous and 3) Others, based on the species’ feeding habits (Ali and Ripley 1973, Jayapal et al 2009). Density estimates were calculated separately for whole strata and the feeding guilds separately. Birds were also classified into ‘habitat guilds’. Each species was classified into three categories 1) Evergreen and Moist Deciduous forest 2) Woodland and 3) Generalist (Ali and Ripley, 1973). Densities were compared using t- tests and one way Analysis of Variance (Zar 1999).

Diversity of birds: The Shannon-Weiner Index was calculated for bird diversity separately for each transect, for both the seasons separately. This index assigns unequal weights to each species based on its frequency of occurrence and is increasingly considered to be the most reliable diversity index (Jost 2007). Hence they served as surrogates for diversity in the habitat attribute analysis.

Software ESTIMATE S 8.0 (Colwell 2006) was used to estimate bird species richness in the same data hierarchy as used in density estimates. The JACK 1 estimator was used in all estimations as it performed consistently with all datasets.

Community similarity

The Morisita- Horn Index and the Sorenson Index were calculated as a measure of community similarity between the three strata in the two strata using software ESTIMATE S 8.0 (Colwell 2006).

Vegetation profile

Structural attributes of the vegetation were calculated using the data from the circular plots.

The following variables were calculated for each transects and then used in the analysis.

- 1) Tree height heterogeneity
- 2) Average girth breast height and its distribution.
- 3) Average canopy cover (quantified for summer and winter separately)
- 4) Tree density
- 5) Epiphytic parasitic abundance/ plot
- 6) Mean N. D. V. I. was extracted from a buffer of 20 m along transects using Arc GIS 9.2 (E.S.R.I. 2003). This was used a measure of canopy continuity.

- 7) Vertical spread of canopy which was calculated as the standard deviation of the difference between the top height and first branch heights of trees.

Floristic attributes

The Shannon Weiner Index was calculated for tree and shrub species for each transect.

Bird- Habitat Interactions

Principal Component Analysis (P.C.A.) (Zar 1999) was carried out for the structural and floristic attributes measured for the three strata along the transect.

Pearson correlations (Zar 1999) were calculated between bird density and diversity measures and habitat variables. Bird species diversity matrices were regressed against vegetation structural variables matrix and a tree species diversity matrix (Mantel test).

The repeated randomization model was used with 1000 runs in all the pairs of matrices used (McCune and Grace 2002). All statistical analyses were done using Statistical Package for Social Sciences (SPSS) version 11.5 (SPSS Inc).

4. Results

One hundred and nineteen species of birds were observed while sampling (Appendix A) and 247 species were seen in the whole duration of the study in the study area. (Appendix B). Winter sampling recorded 89 bird species while 102 species were recorded in summer. This includes 746 detections of birds in winter sampling and 1,171 detections of birds in summer sampling.

4.1 Density of birds

The density of birds was consistently higher in natural forests than in mature or young teak plantations.

Table 4.1 Overall (strata level) density estimates (birds/ha), the values in parentheses denote % coefficient of variance.

Strata	Winter	Summer
Natural Forest	14.95 (9.66%)	22.38 (11.9%)
Mature Plantation	9.94 (10.51%)	15.17 (9.66%)
Young Plantation	10.01 (13.79%)	18.74 (10.88%)

Table 4.2 and 4.3 Guild level density estimates (Birds/ha) for each season - the values in parentheses denote % coefficient of variance

Winter	Natural	Mature	Young
Insectivorous(I) Guild	9.36 (10.41 %)	5.65 (14.12%)	3.90 (18.38%)
Phytophagous (P) Guild	4.20 (28.76%)	3.42 (31.19%)	6.60 (25%)
Others Guild	0.84 (29%)	2.03 (29.44%)	0.83 (24.28%)

Summer	Natural	Mature	Young
Insectivorous (I) Guild	12.06 (10.50%)	6.82 (14.8%)	8.43 (16.58%)
Phytophagous (P) Guild	4.52 (26.7%)	7.65 (28.39%)	10.14 (26.9%)
Others Guild	2.37(32.39%)	2.37 (32.39%)	1.38 (30.68%)

4.1.1 Overall densities

Overall densities at the stratum level were higher in summer than in winter.

Natural forest summer > Natural forest winter ($t_{(0.05, 4, 2)}=2.4517$)

Mature plantation summer > Mature plantation winter ($t_{(0.05, 4, 2)}= 2.8313$)

Young plantation summer > Young plantation winter ($t_{(0.05, 4, 2)}=3.5427$)

4.1.2 Guild level densities amongst strata in the same season-

Insectivorous guild

Winter: Tukey's Post hoc test revealed that natural forests had higher density of birds than young plantations in winter ($F_{(6, 241)}$, $p < 0.05$). Densities of birds in the plantation strata in winter were not significantly different.

Summer- There was no significant difference in insectivorous bird density across the three strata in summer.

Phytophagous guild

Winter- Tukey's Post Hoc test revealed that in winter, Phytophagous guild density in young plantations was higher than mature plantation but not significantly higher than natural forest ($F_{(5, 689)}$, $p < 0.05$).

Summer- Density of phytophagous birds was higher in young plantations than in natural forests and mature plantations ($F_{(14, 302)}$, $p < 0.01$).

Others guild

Winter- Tukey's Post Hoc test showed that the densities of this guild in winter were not significantly different in the three strata.

Summer- The density of this guild in natural forest was significantly different from that in young plantations, but the plantation strata did not differ significantly in their Others guild densities ($F_{(56.61)} p < 0.05$)

4.1.3 Guild level densities across seasons

Insectivorous guild: Insectivorous bird density in natural forests was significantly higher in summer than in winter ($t_{(0.05,4,2)} = 2.94$). Mature plantations did not differ in their densities and the densities in young plantations in summer were higher than the density in winter ($t_{(0.05,4,2)} = 4.9897$).

Phytophagous guild: The density of birds belonging to this guild was higher in mature plantations in summer ($t_{(0.05,4,2)} = 3.0205$). The other two strata did not differ in their Phytophagous guild densities across seasons.

Others guild: The density of this guild was significantly higher in summer than in winter ($t_{(0.05,4,2)} = 5.9294$). The plantation strata did not differ significantly in the density of this guild across seasons.

Proportion of detections in each guild across seasons: The rise in overall density estimates fails to explain the contribution of each guild, an idea can be derived as to the biggest contributor to the rise in density in each stratum by looking at the proportion of densities of each guild across seasons in the same stratum.




Fig 4.1




Fig 4.2




Fig 4.3

Figures 4.1- 4.3

The proportion of each guild in total density in each stratum across seasons.

4.1.4 Comparison of abundances according to habitat guilds across seasons

A comparison of the proportion of detections of each habitat guild in the three strata is a good representation of the change in their abundance. This is because a rise in their density might be an artefact of the overall rise in densities in summer. EvMD shows increase in all three strata with greatest gain in young plantations. While woodland birds do not show a marked change, the overall detections of generalist birds decreases in plantation strata. All detections considered were the ones falling in the first 15m on either side of the transect as detection probability for all size classes at this distance was close to, or equal to 1.




Figure 4.4




Figure 4.5




Figure 4.6

Figures 4.4- 4.6 Graphs showing the proportion of detection of each habitat guild in 15m belt in transects.

4.2 Diversity of birds in natural forests and teak plantations

Estimated species richness of birds showed an increase from winter to summer especially in the plantation strata. The biggest gain in richness occurred in the insectivorous guild in summer.

4.2.1 Winter

Overall estimated bird species richness was highest in natural forests followed by mature plantations and young plantations. At the guild level, insectivorous bird species richness was similar in mature and young plantation strata but less than the natural forests. The phytophagous guild showed a trend of decreasing richness similar to that of overall species richness.

Estimated species richness winter




Figure 4.7 Overall estimated species richness in winter (error bars denote standard deviation)




Figure 4.8 Estimated species richness of the insectivorous guild winter (error bars denote standard deviation)




Figure 4.9 Estimated species richness for phytophagous guild winter (error bars denote standard deviation)

Table 4.4 Estimated species richness in winter using JACK 1 estimator –

	Natural	Mature	Young
Overall	89.33	67.33	55.61
Insectivorous guild	40.17	36.67	35.92
Phytophagous guild	26.67	14.33	11.99

4.2.2 Summer

The overall estimated species richness was highest for natural forest while the two plantation strata showed little difference in richness. At the guild level, the insectivorous guild showed a higher richness in the plantation strata than the natural forest. The phytophagous guild had highest richness in natural forests followed by young plantations and mature plantations.

Estimated Species Richness Summer-




Figure 4.10 The overall estimated species richness in summer (error bars denote standard deviation)




Figure 4.11 The estimated species richness of the insectivorous guild summer (error bars denote standard deviation)




Figure 4.12 The estimated species richness for phytophagous guild summer (error bars denote standard deviation)

Table 4.5 Estimated species richness in summer using the JACK 1 estimator

	Natural	Mature	Young
Overall	102.71	90.29	89.43
Insectivorous	39	42.86	42.71
Phytophagous	29.14	21.87	26

4.2.3 Diversity of birds- Shannon's index calculated at the transect level produced the following index values-

Table 4.6 The Shannon's Index values for each transect in the two seasons

Strata	Transect	Shannon's Index	
		Winter	Summer
Natural	Chipli	2.91	3.58
Natural	Manchale	2.88	3.41
Natural	Konehosur	3.30	3.44
Mature	Ullur	2.58	3.04
Mature	Puradasara	2.83	3.58
Mature	Tupur	3.33	3.12
Young	Tyagarti	2.39	2.67
Young	Jiglemane	2.31	2.34
Young	Konehosur(Y)	2.53	2.89

4.3 Community Similarity

4.3.1 Winter-

Table 4.7 Community similarity using the Morisita Horn Index for winter. Values in parentheses denote number of species shared between the two strata.

Strata	Unique Species			Morisita Horn Index
	Natural	Mature	Young	
Natural-Mature (44)	18	9	-	0.734
Natural-Young (31)	29	-	11	0.624
Mature-Young (31)	-	22	11	0.805

Summer-

Table 4.8 Community similarity using the Morisita Horn Index for summer. Values in parentheses denote number of species shared between the two strata.

Strata	Unique Species			Morisita Horn Index
	Natural	Mature	Young	
Natural-Mature (61)	18	11	-	0.867
Natural - Young (51)	28	-	14	0.797
Mature - Young (53)	-	19	12	0.829




Figure 4.13 Community similarity between sets of strata across seasons (Sorenson's Index)

The three strata show a high degree of bird community similarity. The two plantation strata show highest degree of similarity in winter. In summer however, natural forest and mature plantations show the highest degree of similarity. The overall similarity between the strata increases from winter to summer (high overlap) although the two plantations do not show an increase in similarity. This trend is mirrored in the Sorenson index demonstrating that it is not biased due to many singletons in the plantation strata.

4.4 Vegetation

4.4.1 Structural Attributes

Vertical heterogeneity of a habitat has been recognized as an important factor affecting bird diversity. Foliage height diversity has been quantified as an important vertical heterogeneity variable (Macarthur and Macarthur 1961). The foliage height diversity (F.H.D) of the three strata showed three distinct trends.

Young plantations had a skewed distribution in the lower height categories with most of the foliage located 5-9 m above ground. Mature plantations had a more even foliage distribution with distinct strata forming at 2-5m, 7-9m, and 12-16m above ground. Natural forests had strata at 2-4m and 10-15m above ground.




Figure 4.14 Distribution of foliage in a vertical spread of vegetation

Top Height Heterogeneity

The natural forest had a lot of heterogeneity in top height while the two plantation strata showed more consistent homogenous distribution of tree top height. This was used in further analysis as a measure of the vertical spread and distribution of foliage.




Figure 4.15 Variability among the crown height of trees in each transect. (error bars denote standard deviation)

Vertical spread of vegetation




Figure 4.16 Difference in the vertical spread of foliage in the three strata. (error bars denote standard deviation)

The vertical spread of vegetation was most heterogeneous in natural forests and quite uniform in the plantation strata.

G.B.H: Average G.B.H decreased in the order from natural forests to mature plantations to young plantations. While plantation strata had an even G.B.H. distribution natural forests had a high heterogeneity in its GBH distribution.




Figure 4.17 Distribution of Girth at Breast Height in trees in the three strata.




Figure 4.18 Variability in the GBH in trees of the three strata. (error bars denote standard deviation)

Tree Density




Figure 4.19 Trees/m² in the three strata.

Tree density followed a trend opposite to that of GBH. Tree density was highest in young plantations followed by mature plantations and natural forests (only trees above 15 cm G.B.H were considered)

Epiphytic parasites

Parasites were most abundant in the young plantations as they are known to be partial to areas with high incident sunlight and canopy openings.




Figure 4.20 Number of parasites/ vegetation sampling plot (error bars denote standard deviation)

Canopy cover: Natural forests had almost complete canopy cover in winter, but most of the trees shed their leaves with the onset of summer, the teak had dried leaves in winter and almost no leaves in summer.




Figure 4.21 Seasonal differences in canopy cover in the three strata (error bars denote standard deviation)

4.4.2 Floristic attributes of the three strata

4.4.2.1 Tree diversity: Estimated tree species richness was highest in natural forests followed by mature plantations and young plantations.




Figure 4.22 Estimated tree species richness using the JACK 1 estimator (error bars denote standard deviation)

4.4.2.2 Tree and shrub species diversity at the transect level

Table 4.9 The Shannon's index for tree and shrub diversity for transects

	Transect	Tree	Shrub
Natural	Chipli	1.09	1.72
	Manchale	1.88	1.05
	Konehosur	2	1.08
Mature	Ullur	1.39	0.09
	Puradasara	1.8	0.03
	Tupur	1.39	0.07
Young	Tygarti	0.96	1.84
	Jiglemane	1.17	1.97
	Konehosur Young	1.08	0.96

Natural forests: These are habitats with low tree density, a highly heterogeneous GBH distribution, highest heterogeneity in tree height diversity ($F_{(10,801)} p < 0.05$) and vertical spread of foliage ($F_{(7,345)} p < 0.05$). It had high tree and shrub diversity with a high and continuous canopy cover and few epiphytic parasites.

Mature plantations: As these are actively managed habitats they have homogenous GBH distribution, tree density is slightly higher than natural forests and a homogenous tree height distribution. Canopy cover was highly variable as teak is a deciduous species. Epiphytic parasites were a little more abundant than in natural forests. There is an abundant regeneration of trees like *Xylia xylocarpa*, *Terminalia paniculata* and *Syzigium cumini*. Tree species diversity is only a little below that of natural forests. But they were very poor in shrub species diversity.

Young plantations: These habitats have a high tree density, and highly homogenous GBH distribution with most trees between 20cm -50cm in girth and homogeneity in tree height. They have highly variable and discontinuous canopy cover. Teak trees in young plantations lost their leaves earlier and grew new leaves earlier than the mature plantations (pers. observation). They had the highest density of epiphytic parasites ($F_{(8,691)} p < 0.05$). Tree diversity was not very high but they had the highest shrub species diversity ($F_{(13,029)} p < 0.05$).

4.5 Habitat correlates explaining bird diversity

The Principal Component Analysis decomposed the data into three principal factors together explaining 90.28 % of the variance in the dataset.

Table 4.10 Contribution of each variable to the extracted factors

Component Matrix

% Variance explained	Component		
	(1) 48.92	(2) 25.49	(3) 15.87
Canopy Cover (Summer)	.419	.491	.735*
GBH	.824*	-.418	.201
Std.dev of Top Height Diversity	.969*	-.044	-.190
Tree Species Diversity	.599	.728*	.103
Shrub Species Diversity	-.267	-.601	.678
Tree Density	-.676	.686*	.080
Shrub Density	.862*	.056	-.134

Extraction Method: Principal Component Analysis.

a 3 components extracted.

(*) indicate significant correlation between Factor Score and Variable.

Component 1 was related to “forest maturity” as GBH, Top Height Diversity and Shrub Density were all contributing to this factor. Component 2 was related to tree density and diversity, while Component 3 was related to canopy cover.

4.5.2 Habitat variables explaining trends in bird density and diversity

Winter

Overall bird diversity was explained by N.D.V.I. and canopy cover ($R= 0.737$ $p< 0.01$ and $R=0.807$ $p<0.05$ respectively). This shows that birds favour areas with high canopy cover and respond positively to canopy connectivity and continuity. It also shows a strong positive correlation to the tree height heterogeneity ($R= 0.871$, $p<$

0.01). Hence bird diversity increases with increasing heterogeneity in vertical spread of foliage. The insectivorous guild density also responded to tree height heterogeneity ($R= 0.793$ $p < 0.01$). Phytophagous birds showed a positive correlation to mean number of epiphytic parasites/ plot ($R= 0.766$, $p < 0.05$). These parasites were flowering in winter and might have attracted a large number of nectarivores. They also showed a negative correlation to tree height heterogeneity ($R= 0.815$, $p < 0.01$) which could be an artefact of them being most abundant in young plantations.

Overall resident bird composition and tree species composition showed a strong relationship ($R= 0.49$, $p < 0.001$). Resident insectivorous and phytophagous guilds also showed significant relationships ($R= 0.36$ $p < 0.001$, $R= 0.49$, $p < 0.001$ respectively) (Mantel tests). This shows that tree composition plays a significant role in determining the resident bird community.

Summer-

In summer overall bird diversity was highly positively correlated to Principal Component 1 which represented forest maturity ($R= 0.852$, $p < 0.01$). Overall diversity also shows a strong correlation to the tree height heterogeneity ($R= 0.820$, $p < 0.01$). Insectivorous guild density was negatively correlated to tree density ($R= -0.824$, $p < 0.01$) and positively correlated to G.B.H ($R= 0.864$, $p < 0.05$) suggesting that this guild responded well to forest maturity. Phytophagous bird density was positively correlated to number of epiphytic parasites ($R= 0.796$, $p < 0.01$) and negatively to tree density and diversity ($R= -0.885$, $p < 0.01$) due to highest densities of this guild occurring in young plantations. This guild is negatively correlated to the tree height heterogeneity ($R= -0.815$, $p < 0.01$). This is because young plantations had the least vertical heterogeneity in habitat and had highest densities of this guild.

There was no significant correlation between bird composition and structural habitat variables and tree species composition (Mantel tests).

5. Discussion

Most studies have shown that forest plantations support less avifaunal diversity than natural forests. However the differences in seasonal use of plantations have not been discussed widely. This study demonstrates that a well known trend of decreasing bird diversity and abundance from natural forests to mature plantations to young plantations exists in teak plantations also. However the results indicate a distinct change in both bird density and diversity with change in season.

5.1 Differences in Habitat Characteristics (Vegetation structure)

Teak plantations of Sagar Forest Division show some obvious structural and floristic differences from natural forests. As they are actively managed forests, teak plantations show an increased tree density and lowered tree species diversity. Floristically it is difficult to tease out differences in tree and shrub diversity, as they do not show a high regeneration of teak saplings. *Xylia xylocarpa* *Terminalia paniculata* and *Diospyros melanoxylon* are the dominant regenerating species in all the strata. This fact may contribute to the similarity in the shrub diversity. Teak plantations also have a discontinuous canopy because teak trees as a physiognomic trait do not have a high horizontal canopy spread. They also show a high degree of seasonal change in canopy cover as almost all the trees shed leaves simultaneously resulting in a considerable amount of insolation (incident sunlight). This has resulted in young teak plantations having the highest shrub diversity due to abundant light reaching the forest floor. An important feature of young plantations is the increased abundance of epiphytic parasites on the teak trees. Additionally all teak plantations have an increased abundance of the climber *Calycopteris floribunda*, a dry deciduous plant utilizing the abundant sunlight. However, the most important structural attribute that sets natural forests apart from the actively managed teak plantations is the

heterogeneity in the top heights of trees and the vertical spread of foliage rendering the natural forests much more structurally complex than the plantation strata.

5.2 Density of birds

The increase in over all densities from winter to summer is a phenomenon seen often with birds in forest environments (Karr 1976). This season is the breeding season for most birds and they might be consequentially converging into forest habitats for nesting. Findings of this study show that there is ingression of individuals of species that were common to both seasons into forest habitats in summer. Number of detections of species common to both seasons increase in summer. Subtraction of winter values from those in summer give positive values (an increase in abundance) for around 65 % species in all strata. The species showing negative values are primarily migratory birds like Greenisg Warbler (*Phylloscopus trochiloides*) and Western Crowned Warbler (*Phylloscopus occipitalis*).




Figure 5.1 Increase in abundance of birds common to both seasons in natural forests. A positive value bar denotes an increase in abundance in summer.




Figure 5.2 Increase in abundance of birds common to both seasons in mature plantations. A positive value bar denotes an increase in abundance in summer.




Figure 5.3 Increase in abundance of birds common to both seasons in young plantations. A positive value bar denotes an increase in abundance in summer.

A look at the ecology of the species increasing in abundance throws some light on the possible reasons for the same. The species that shows the highest increase across the three strata is the White rumped shama (*Copsychus malabaricus*) which is a cavity nesting bird (Ali and Ripley 1973). Even the other species increasing in abundance are mostly birds that are cavity nesters like the Greater flameback (*Chrysocolaptes lucidus*) Black rumped flameback (*Dinopium bengalense*) and Brown headed barbet (*Megalaima zelanica*). They also include birds that need a good shrub layer as they nest close to the ground like Yellow browed bulbul (*Iole indica*), Red whiskered bulbul (*Pycnonotus jocosus*), Brown cheeked fulvetta (*Alcippe poioicephala*), Indian scimitar babbler (*Pomatorhynus horsefieldii*) and Tickell's blue flycatcher

(*Muscicapa tickellii*) (Ali and Ripley 1973). There is a strong possibility that the birds use plantations as nesting habitats, also possibly due to these habitats have minimal disturbance due to enforced protection.

Overall density of birds was highest in natural forests. Prachi Mehta (1998) in her study in the Bori Wildlife Sanctaury reports that the density of birds increases from young teak plantations to mature teak plantations with maximum density in natural forests, a trend published for other plantations as well (Trivedi, 2006 Rumble and Gobeille, 1998). A similar trend is also reflected in the current study also and this is attributed generally to the stabilization of habitat structure a more stable microclimate regime and plant communities. Guild level densities of birds however showed some peculiar results. The plantation strata show a high variability in bird abundance.

While the density of insectivorous birds did not change in natural forests across seasons, plantations showed a marked increase in insectivore density in summer. This might be a response to the increased arthropod abundance that occurs in most forest environments in summer (Karr 1976) and also an apparent tracking of arthropod abundance after the teak starts getting new leaves.

Phytophagous birds show a reverse trend in density. In both seasons young plantations supported the highest densities of phytophagous birds. This appears to be primarily explained by the the high abundance of Common rosefinch (*Carpodacus erythrinus*). These birds were found in large flocks in the young plantations. Bamboo (*Dendrocalamus strictus*) was flowering and holding seed in most of the study area. As open forest granivorous birds, Common rosefinches probably preferred young plantations to the closed moist-deciduous and semi-evergreen natural forests even though they did not have the highest bamboo density. Common rosefinches make up a

large portion of the Phytophagous guild detections. They are followed in abundance by sunbirds and flowerpeckers. These birds were probably attracted to the epiphytic parasites that were found in highest densities in this stratum. These parasites were flowering during winter and fruiting during summer, a constant food source explaining the inflated abundance of these birds. Young plantations supported no true frugivores in winter and very few in summer.




Figure 5.4 Rank abundance of phytophagous birds in young plantations in winter.




Figure 5.5 Rank abundance of phytophagous birds in young plantations in summer.

Hence the high density of phytophagous birds was mostly due to the inflated abundance of nectarivores and rosefinches. This abundance of nectarivores might have an explanation in the resource concentration hypothesis. This phenomenon has been observed with other taxa like insects in monocultures (Connors et al 2000). This finding is in contrast to that of Bell (2000) who reported a loss of granivores and nectarivores from teak plantations in New Guinea where it is an exotic and the local epiphytic plants are probably not adapted to exploit teak.

5.3 Diversity

The diversity of birds also followed a similar trend as that of density with natural forests showing highest diversity followed by mature plantations and then by young plantations in winter. However this difference is not apparent in summer. Bhagwat et al (2008) report that on an average only 50 - 60% of original bird fauna is represented in forest plantations. Generally the birds that are missing from plantations are

primarily the obligate frugivores and birds of primary closed canopy forest (Petit et al 1999, Raman 2001, Daily et al 2001). Hence plantations of native species, even though they support comparable species richness, generally do not contain species of conservation importance (Kwok and Corlett 2000, Rotenburg 2007). The present study looks at seasonal variation in diversity and variation in the community.

The increased bird diversity in all three strata in summer can be attributed to the influx of species for nesting (pers. observ.) or for feeding opportunities. Interestingly the species richness of even young plantations increases considerably and is as high as the mature plantations. The greatest rise in estimated species richness which is amongst phytophagous birds is the effect of many species like Oriental turtle doves (*Streptopelia orientalis*), Vernal hanging parrots (*Loriculus vernalis*), White rumped munias (*Lonchura striata*) and Chestnut shouldered petronia (*Petronia xanthocollis*) using the plantations to feed on the fruits of *Calycopteris floribunda* a common climber which has a large fruit crop.

In part, a common influence on the increase in density and diversity is the apparent seasonal use of the plantation habitats by evergreen and moist deciduous forest (EvMD) birds. The birds belonging to this habitat guild show a consistent rise across all strata, with the highest gain in young plantations. These birds include species like Yellow browed bulbul (*Iole indica*), White bellied woodpecker (*Dryocopus javensis*), Brown-cheeked fulvetta (*Alcippe poiocephala*), Large woodshrike (*Tephrodornis gularis*), Malabar Trogon (*Harpactes fasciatus*), Dark fronted babbler (*Rhopocichla atriceps*) and Pompadour green pigeons (*Treron pompadora*). The reason for this seasonal influx might be for nesting (pers. observation) or to exploit feeding opportunities. In their study of birds of Uttara Kannada Daniels et al (1992) said that birds in teak plantations have species in common with dry deciduous forests. The

present study also supports this hypothesis due to the high proportion of woodland and generalist species, but also reports a possibility of interplay of seasonality and use of these plantations by evergreen forest elements in summer.

5.4 Community Structure

The bird community of the region shows a high degree of overlap. The similarity increases in summer likely due to the influx of closed forest birds into the plantation strata. This is supported by both the Morisita Horn Index and the Sorenson Index indicating that the high degree of similarity is not an artefact of individual species abundances.

Overall trends in bird abundance and diversity

Winter

Density

Overall	Natural forests > Mature plantations \approx Young plantations
Insectivore guild	Natural forests > Mature plantations \approx Young plantations
Phytophagous guild	Natural forests < Mature plantations < Young plantations

Diversity

Overall	Natural forests > Mature plantations > Young plantations
Insectivore guild	Natural forests > Mature plantations > Young plantations
Phytophagous guild	Natural forests > Mature plantations > Young plantations

Summer

Density

Overall	Natural forests > Young plantations > Mature plantations
Insectivore guild	Natural forests > Young plantations > Mature plantations
Phytophagous guild	Natural forests < Mature plantations < Young plantations

Diversity

Overall	Natural forests > Mature plantations \approx Young plantations
Insectivore guild	Natural forests \approx Mature plantations \approx Young plantations
Phytophagous guild	Natural forests \approx Mature plantations \approx Young plantations

Alpha, beta and gamma diversity are amongst the fundamental descriptive variables of ecology, but their quantitative definition has been controversial. The dependence of the value of beta on alpha has been seen as a cause of concern, as it makes the diversity index behave in ways that are not always intuitively correct. The numbers equivalent of all standard diversity indices however behave in this intuitive manner. This has led to the modification of Shannon Weiner Index to give the numbers equivalent of the formula as $[\exp \{-\sum p_i * \log(p_i)\}]$. By this new modification, the landscape level diversity is $\exp(H_\gamma) = \exp(H_\alpha) + \exp(H_\beta)$ i.e. $\exp(H_\beta) = \exp(H_\gamma) - \exp(H_\alpha)$.

If the numbers equivalent answer for this equation is ≥ 2 then it is safe to say that the samples are drawn from two mutually exclusive and distinct communities (Jost 2007). Gamma level diversity was calculated for the whole dataset for the two seasons separately. Then, using the diversity of the natural forest strata as the alpha diversity (as it had the highest species richness) the beta diversity was calculated. It was 1.12 for winter and 1.21 for summer. Hence it seems that the bird community of the teak plantations of the area is derived from the original Semi Evergreen- Moist Deciduous forest bird community.

5.5 Habitat Attributes

Forest structure (physiognomy) and composition (floristics) have been known to be the two most important albeit broad, factors governing the avian community of a given area. Also there is evidence that these factors play important roles at different levels in the landscape. While floristics play a role at the site level, forest structure often plays a role at the regional level (Lee and Rotenberry 2005). Hence vegetation structure plays a role across habitats and vegetation composition plays a role within a habitat. Similar trends have been seen in the present study. As comparisons were done

across three main habitats, vegetation structure proved to be the most important factor explaining differences in bird density and diversity.

5.5.1 Patterns in bird responses common to both seasons

An important factor determining bird density and diversity was the vertical structural heterogeneity (tree height heterogeneity). This attribute renders vertical structural complexity which provides for more niches to develop and hence results in higher bird densities (MacArthur and MacArthur 1961, Holmes et al 1986, Mitra and Sheldon 1993, Hansen et al 1995). Insectivores in both seasons responded to high basal area which formed a structural gradient separating the three strata ($F_{(5,160)} p < 0.05$), that was greatest in natural forests. This resulted in a high density and diversity of this guild in these habitats. Phytophagous guild densities were primarily governed by epiphytic parasite abundance and all the other correlations, like the relationship with high tree density and low GBH and low canopy cover are an artefact of the same.

5.5.2 Seasonal Patterns

Winter

Contrary to Daniels et al (1992) who reported that bird species diversity increased with increased patchiness of canopy, the current study shows a strong correlation of overall winter bird diversity and canopy intactness and continuity. In fact canopy cover also explains the density of insectivorous guild in winter. Hence canopy cover and intactness are the most important governing factors of bird diversity in the winter.

Canopy cover has been recognized as an important factor for bird diversity in many studies (Mazerrole and Villard 1999, Rotenburg 2007, Raman 1998, Wilson 1974, Malcom et al 2006). Most of these studies also show that bird diversity of evergreen

and semi-evergreen forests are most severely affected changes in canopy structure, a trend seen in the current study. Resident bird composition was also correlated to tree composition a plausible phenomenon as natural forests with high floristic diversity offers food resources during the winter 'resource-crunch' period.

In summer, canopy did not play a role in determining bird responses to habitat (as most tree species were leaf less or sprouting new leaves). Although the effect of floristics between habitats is known to be minimal, plant species diversity increases the strength of the correlation between bird diversity and vegetation structure. Tree composition was not correlated to bird composition in summer.

Teak plantations hence emerge as seasonally variable bird habitats. Natural forests being stable ecosystems change very little with seasons in terms of diversity. However teak plantations, although depauperate in terms of diversity and abundance in winters seem to offer subsistence to a large variety of birds with diverse life history traits in summer. Mature plantations are closer to natural forests in terms of structure and floristics and hence are more stable and change minimally across seasons. Young plantations are more dynamic. With their abundant parasites, they are home to an almost unnatural abundance of nectarivores. Insectivores also seem to seasonally exploit their resources to a greater degree than even mature plantations even though they are the most structurally and floristically homogenous of the three strata. This might have some explanation with shrub diversity which is highest in young plantations and almost negligible in mature plantations. Also, the presence of many insectivorous and phytophagous closed forest species within teak plantations during some seasons suggests that closed forest birds may use plantations more than previously believed.

5.6 Conclusion

Teak plantations do not harbour the same diversity and abundance of birds as natural forests especially that of closed forest frugivores and canopy specialist birds. A certain proportion of the landscape has to have native primary forest (possibly in a matrix) so as to function as a species pool. However, some pertinent points emerge as to what might be the characteristics of bird-friendly teak plantations. Distance to natural forests is an obvious and intuitive factor however that variable was controlled for in this study, so it cannot be discussed here. Teak plantations near natural forests may support many native bird species if certain habitat attributes are maintained. Canopy being an important determinant, some trees with large canopy spread like the locally occurring *Dilena* spp. or *Ficus* spp. can be maintained in a mosaic to enhance the canopy structure of the forest. Also certain conventional practices like climber cutting and parasite removal can be less rigorous as these plants seem to enhance bird diversity especially that of phytophagous birds. Also, clear felling which is a common practice for extraction in teak plantations might be replaced by a system in which trees are felled selectively. This will increase the vertical heterogeneity and stratification of the habitat increasing the diversity of birds using the area. Teak plantations being actively guarded by authorities have very little anthropogenic disturbance. The natural forests of the area are heavily exploited for firewood and leaf litter collection in early summer, the breeding season for most bird species causing a great deal of disturbance. Hence the protected teak plantations might also be functioning as good and safe nesting habitats for birds. Also as they are protected from fire, they can grow a healthy shrub under storey which is an important structural component with regards to avifauna commercial gain derived from teak plantations.

Planted forests are neither inherently good nor bad; rather it is the choices we make about how to use them that determine whether they contribute to, or detract from, broader societal goals such as nature conservation. Whether we like it or not, plantations are here to stay and are capable of delivering ancillary social and environmental co-benefits. Dismissing plantations as just another agricultural crop is therefore counterproductive. Through intelligent research and management, we can work towards a system that facilitates the realization of many ancillary goals of plantations while maximizing output of the desired goods.

References

- Ali S., Ripley D.S. (1973) Handbook of the Birds of India and Pakistan, Oxford University Press, New Delhi.
- Anand M.O., Krishnaswamy J, Das A. (2008) Proximity To Forests Drives Bird Conservation Value Of Coffee Plantations: Implications For Certification., *Ecological Applications* 18(7) 1754-1763.
- ANON (2005), State of Forest Report 2005, Chapters 3, 5 and 6, 23-40, Forest Survey of India, Dehradun, India
- ANON (2002), Forest Plantations Working Papers, Forestry Department, Food and Agriculture Organization of the United Nations, Rome (Italy).
- Arnold G (2003) Bird species richness and abundance in wandoo woodland and in tree plantations on farmland at Baker's Hill, Western Australia, *Emu* 103(3) 259 - 269
- Bell H.L. (2000) The Effects on Rain-Forest Birds of Plantings of Teak, *Tectona grandis*, in Papua New Guinea., *Australian Wildlife Research* 6(3) 305 – 318
- Bhagwat S.A., Willis K.J., Birks J.B., Whittaker R.J. (2008), Agroforestry: a refuge for biodiversity, *Trends in Ecology and Evolution* Vol 23(5) 261-267
- Champion H.G., Seth S.K. (1968), Forest Types of India, The Manager of Publications, Delhi.
- Chazdon R. (2008), Beyond Deforestation: Restoring Forests and Ecosystem Services on Degraded Lands, *Science* 320 (5882) 1458-1460

Clout M.N., Gaze P.D. (1984), Effects of plantation forestry on birds in New Zealand,
The Journal of Applied Ecology Vol 21 (3) 795-815

Colwell R.K. (2006) *Estimate S*. Statistical estimation of species richness and shared
species from samples. Version 8. Persistent URL <purl.oclc.org/estimates>

Connors E.F., Courtney A.C. Yoder J.M (2000) Individuals-Area Relationships: The
Relationship between Animal Population Density and Area, Ecology 81 (3)
734-748

Daily G , Ehrlich P, Sanchez-Azofeifa A. (2001) Countryside Biogeography Use of
human-dominated habitats by the avifauna of southern Costa Rica, Ecological
Application 11 (1) 1-13

Daniels R.J.(1989), A Conservation Strategy for the Birds of the Uttara Kannada
District, Phd. Thesis, Centre for Ecological Sciences, Bengaluru.

Daniels R.J., Joshi N.V., Gadgil M. (1992) On The Relationship Between Bird And
Woody Plant Species Diversity In The Uttara Kannada District Of South
India, Ecology 5311-5315

Enters (2000), Site, technology and productivity of teak plantations in south-east Asia,
Unasylva 201 (51) 55-61.

Farwig N., Sajita N, Bohning-Gaese K. (2008) Conservation value of Forest
Plantations on Bird Communities in Western Kenya, Forest Ecology and
Management (in press)

Goodwin J, Fahrig L. (2002) How does landscape structure influence landscape
connectivity?, Oikos 99: 552-570,

- Hansen A, McComb W, Vega R., Raphael M. Hunter M (1995), Bird Habitat Relationships in Natural and Managed Forests in the West Cascades of Oregon, *Ecological Applications*, Vol 5 (3) 555-569
- Harikrishnan S., Mathur P.K., Raphael M.G., Udhayan A., Vasudevan K. (2008), Development of adaptive management approach for phasing out monoculture plantations in the Anamalai Conservation Area, Unpublished report, Management of Forests in India for Biological Diversity and Forest (Phase II).
- Hawrot R, Niemi G (1996) Effects of edge type and patch shape on an avian communities in a mixed conifer-hardwood forest. *The Auk* 113(3):586-598.
- Jende O. Pohlan H, Janssens H.(2005) Conversion process and reforestation practices of coffee plantations in Chiapas, Mexico, Conference on International Agricultural Research for Development, Tropentag, Germany.
- Jayapal R. Qureshi Q. Chellam R.(2007) Developing a Spatial Conservation Protocol for the Central Indian Highlands through a biogeographical analysis of birds and existing Protected Area network: A Geographical Information Systems Approach, Wildlife Institute of India, Dehradun.
- Jayapal R. Qureshi Q. Chellam R. (2009) Importance of forest structure versus floristics to composition of avian assemblages in tropical deciduous forests of Central Highlands, India, *Forest Ecology and Management*, In Press.
- Jost L (2007) Partitioning Diversity into Independent Alpha and Beta Components, *Ecology* 88(10) 2427-2429.

- Karr J. (1976) Seasonality, Resource Availability and Community Diversity in Tropical Bird Communities, *The American Naturalist* 110 (976) 973-994.
- Kwok H.K., Corlett R.T. (2000), The bird communities of a natural secondary forest and a *Lophostemon confertus* plantation in Hong Kong, South China, *Forest Ecology and Management* 130 (1-3) 227-234.
- Lee P.Y., Rottenbury John T, (2005), Relationships between bird species and tree species assemblages in forested habitats of eastern North America, *Journal of Biogeography*, 32 1139-1150.
- Lal J.B.(1989) India's Forests Myth and Reality, Historical Background (pp 15-36), *Silvicultural Principles and Practices* 106-132, Natraj publishers, Dehradun, India.
- MacArthur R.H., MacArthur J.W. (1961) On Bird Species Diversity, *Ecology* 42(3) 592-598
- Marsden S, Whiffin M, Galetti M (2001) Bird diversity and abundance in forest fragments and Eucalyptus plantations around an Atlantic forest reserve, Brazil, *Biodiversity and Conservation* 10 737-751
- Mayeaux P., Holmgren P. Achard F., Hugh E. Stibig H, Branthomme A (2005), Tropical Cover Changes in the 1990s and options for future monitoring, *Philosophical Transactions of the Royal Society B*, **360** 373-384.
- McCune B, Grace J.B. (2002) Mantel tests, *Analysis of Ecological Communities*, MjM software design, Oregon, USA,- 211-217

- Mehta P. (1998), The effects of forestry Practices on bird species diversity in Satpuda hill ranges, Phd. Thesis, Wildlife Institute of India, Dehradun.
- Mitra S.S., Sheldon F.H. (1993) Use of an Exotic Tree Plantation by Bornean Lowland Forest Birds, *The Auk* 110 (3) 529-540
- Moguel P, Toledo V (1999) Review: Biodiversity Conservation in Traditional Coffee Systems of Mexico, *Conservation Biology* 13(1) 11-21
- Mohan D. (2007) Habitat Selection of Birds in New Forest, Dehradun, India. Phd Thesis, Forest Research Institute, Dehradun, India.
- Montagnini F, Cusack D., Petit B., Kanninen M (2005) Environmental Services of Native Tree Plantations and Agroforestry Systems in Central America, The Haworth Press, Inc.
- Murthy A. (2001), Working plan of the Sagar Forest Division, Chapters I-V, Karnataka Forest Department.
- Nagendra H. (2007), Drivers of reforestation in human-dominated forests, *PNAS*, 39 (104) 15218-15223.
- Nair V., Jayson E.A. (1988) Habitat Utilization by Large Mammals in Teak Plantations and Natural Forests, KFRI Research Report, Kerala Forest Research Institute.
- Petit, L.J., Petit, D.R., Christian, D.G., Powell, H.D.W (1999) Bird communities of natural and modified habitats in Panama. *Ecography* 22 292-304.

- Raman, T.R.S. (2001). Effect of slash-and-burn shifting cultivation on rainforest birds in Mizoram, northeast India, *Conservation Biology* 15 685– 698.
- Raman T.R.S (2006), Effects of habitat structure and adjacent habitats on birds in tropical rainforest fragments and shaded plantations in the Western Ghats, India, *Biodiversity and Conservation* 15 1577-1607.
- Rosenstock, S. S., D. R. Anderson, K. M. Giesen, T. Leukering, and M. F. Carter.(2002) Landbird counting techniques: current practices and an alternative. *The Auk*, **119**, 46-53.
- Rotenburg J (2007) Ecological role of a tree (*gmelina arborea*) plantation in guatemala: an assessment of an alternative land use for tropical avian conservation, *The Auk* 94 (2) 201-210
- Rumble M, Gobeille J (1998) Bird Community Relationships to Succession in Green Ash (*Fraxinus pennsylvanica*) Woodlands, *American Midland Naturalist*, Vol 140 (2) 372-381.
- Saha (2001) Vegetation composition and structure of *Tectona grandis* (teak, Family Verbanaceae) plantations and dry deciduous forests in central India Forest Ecology and Management Volume 148, (1-3), 159-167
- Sekercioglu C.H.(2002) Effects of forestry practices on vegetation structure and bird community of Kibale National Park, Uganda, *Biological Conservation* 107 229-240

- Shochat E, Abramsky Z, Pinshow B (2001) Breeding Bird Species Diversity in the Negev: Effects of Scrub Fragmentation by Planted Forests, *Journal of Applied Ecology*, 38 (5) 1137-1147
- Tejeda-Cruz C, Sutherland W.J. (2006) Bird responses to shade coffee plantations, *Animal Conservation*, Vol 7 (2) 169-179.
- Telleria, J.L., Santos T., Sanchez A., Galarza A., (1992) Habitat structure predicts bird diversity distribution in Iberian forests better than climate, *Bird Study* Vol. 39(1), 63-68.
- Thomas L., Buckland S.T., Burnham K.P., Anderson R.D., Jeffrey L. Laake, Borchers D.L., Strindberg S. (2002), *Distance sampling*, Volume 1 544–552, *Encyclopedia of Environmetrics*, John Wiley & Sons, Ltd, Chichester.
- Thomas L., Buckland S.T., Burnham K.P., Anderson R.D., Jeffrey L. Laake, Borchers D.L., Strindberg S. (2006) *DISTANCE* version 5, Research Unit for Wildlife Population Assessment, University of St. Andrews, United Kingdom.
- Trivedi P. (2006), *Ecology and Conservation of Avifauna of Some Forested Areas in Gujarat, India*, Unpublished Phd. Thesis, Saurashtra University.
- Verner J.(1985), *Assesment of Counting Techniques*, *Current Ornithology* Volume 2, Chapter 8 247-304.
- Wethered R, Lawes M.J. (2005) Nestedness Of Bird Assemblages In Fragmented Afromontane Forest: The Effect Of Plantation Forestry In The Matrix, *Biological Conservation* 123 125-137

Wilson M.F.(1974) Avian Community Organization and Habitat Structure, *Ecology*
55(5), 1017-1029.

Wright J., Muller-Landau H(2006), The Future of Tropical Forest Species, *Biotropica*
38(3): 287–301

Zar J.H. (1999), *Biostatistical Analysis*, Pearson Education, Prentice Hall New Jersey.

APPENDIX A -Bird species used in the analysis

Common Name	Scientific Name	Habitat Guild	Feeding Guild	Seen in
Ashy Drongo	<i>Dicrurus leucophaeus</i>	G	I	W S
Ashy Prinia	<i>Prinia socialis</i>	G	I	S
Asian Brown Flycatcher	<i>Muscicapa dauurica</i>	G	I	S
Banded bay Cuckoo	<i>Cacomantis sonneratii</i>	G	O	S
Black crested Bulbul	<i>Pycnonotus melanicterus</i>	EvMD	O	S
Black headed Cuckooshrike	<i>Coracina melanoptera</i>	W	I	S
Black hooded Oriole	<i>Oriolus xanthornus</i>	G	O	W S
Black naped Monarch	<i>Hypothymis azurea</i>	G	I	W S
Black naped Oriole	<i>Oriolus chinensis</i>	Ev MD	O	S
Black rumped Flameback	<i>Dinopium benghalense</i>	G	I	W S
Blue capped Rock thrush	<i>Monticola cinclorhynchus</i>	Ev mD	O	S
Blue faced Malkoha	<i>Phaenicophaeus viridirostris</i>	Ev MD	O	S
Blue tailed Beeater	<i>Merops philippinus</i>	G	I	S
Blyth's Reed Wabler	<i>Acrocephalus dumetorum</i>	G	I	S
Booted Warbler	<i>Hippolais caligata</i>	G	I	W
Bronzed Dongo	<i>Dicrurus aeneus</i>	Ev MD	I	W S
Brown capped pygmy Woodpecker	<i>Dendrocopos moluccensis</i>	W	I	W S
Brown headed Barbet	<i>Megalaima zeylanica</i>	G	P	W S
Brown Shrike	<i>Lanius cristatus</i>	W	I	W
Chestnut shouldered Petronia	<i>Petronia xanthocollis</i>	G	P	S
Chestnut tailed Starling	<i>Sturnia malabarica</i>	W	O	S
Common Rosefinch	<i>Carpodacus erythrinus</i>	G	P	W S
Common Drongo	<i>Dicrurus macrocercus</i>	G	I	W S
Common Hawk Cuckoo	<i>Cuculus varius</i>	W	I	S
Common Myna	<i>Acridotheres tristis</i>	G	O	S
Common Woodshrike	<i>Tephrodornis pondicerianus</i>	G	I	S
Coppersmith Barbet	<i>Megalaima haemacephala</i>	G	P	W S
Coucal	<i>Centropus sinensis</i>	G	O	W S
Crimson backed Sunbird	<i>Leptocoma minima</i>	Ev MD	P	W S
Crimson Fronted Barbet	<i>Megalaima rubricapillus</i>	Ev MD	P	S
Crimson Sunbird	<i>Aethopyga vigorsii</i>	G	P	S
Dark fonted Babbler	<i>Rhopocichla atriceps</i>	Ev MD	I	W S
Emerald Dove	<i>Chalcophaps indica</i>	Ev MD	P	W S
Eurasian Blackbird	<i>Turdus merula</i>	Ev MD	O	W
Fairy Bluebird	<i>Irena puella</i>	Ev MD	O	S
Forest Wagtail	<i>Dendronanthus indicus</i>	G	I	W S
Gold fronted Chroloopsis	<i>Chloropsis aurifrons</i>	W	O	W S
Golden Oriole	<i>Oriolus oriolus</i>	G	O	W S
Great Tit	<i>Parus major</i>	G	O	W
Greater Flameback	<i>Chrysocolaptes lucidus</i>	Ev MD	I	W S
Greenish Warbler	<i>Phylloscopus trochiloides</i>	G	I	W S
Grey headed Flycatcher	<i>Culicicapa ceylonensis</i>	G	I	W

Hearspotted Woodpecker	<i>Hemicircus canente</i>	Ev MD	I	W S
Hume's Warbler	<i>Phylloscopus humei</i>	G	I	S
Indian Pitta	<i>Pitta brachyura</i>	G	O	S
Indian Scimitar Babbler	<i>Pomatorhinus horsfieldii</i>	G	I	W S
Iora	<i>Aegithina tiphia</i>	G	I	W S
Jungle Babbler	<i>Turdoides striata</i>	G	I	W S
Jungle Crow	<i>Corvus macrorhynchos</i>	G	O	W S
Jungle Myna	<i>Acridotheres fuscus</i>	W	O	S
Koel	<i>Eudynamys scolopaceus</i>	G	O	W S
Large Cuckooshrike	<i>Coracina macei</i>	G	O	W S
Large Woodhsrike	<i>Tephrodornis gularis</i>	Ev MD	I	W S
Lesser Yellownape	<i>Picus chlorolophus</i>	Ev MD	I	S
Loten's Sunbird	<i>Cinnyris lotenius</i>	G	P	S
Magpie Robin	<i>Copsychus saularis</i>	G	I	W S
Malabar Grey Hornbill	<i>Ocyeros griseus</i>	Ev MD	O	W S
Malabar Parakeet	<i>Psittacula columboides</i>	Ev MD	P	S
Malabar Pied Hornbill	<i>Anthracoseros coronatus</i>	Ev MD	P	S
Malabar Trogon	<i>Harpactes fasciatus</i>	Ev MD	O	W S
Malabar whistling thrush	<i>Myophonus horsfieldii</i>	EvMD	O	W
Nilgiri Flycatcher	<i>Eumyias albicaudatus</i>	Ev MD	I	S
Olive backed pipit	<i>Anthus hodgsoni</i>	G	I	S
Orange headed Thrush	<i>Zoothera citrina</i>	G	O	W S
Oriental Turtle Dove	<i>Streptopelia orientalis</i>	Ev MD	P	W S
Oriental White eye	<i>Zosterops palpebrosus</i>	G	O	W S
Pale billed Floerpecker	<i>Dicaeum erythrorhynchos</i>	G	P	W S
Paradise Flyatcher	<i>Terpsiphone paradisi</i>	G	I	W S
Pied Flycatchershrike	<i>Hemipus picatus</i>	G	I	W S
Plain Flowepecker	<i>Dicaeum concolor</i>	Ev MD	P	W S
Plum headed Parakeet	<i>Psittacula cyanocephala</i>	G	P	W S
Pompadour Green Pigeon	<i>Treron pompadora</i>	EvMD	P	W S
Puff Throated Babbler	<i>Pellorneum ruficeps</i>	G	I	S
Purple rumped Sunbird	<i>Leptocoma zeylonica</i>	G	P	W S
Purple Sunbird	<i>Cinnyris asiaticus</i>	G	P	W S
Quaker Babbler	<i>Alcippe poioicephala</i>	Ev MD	I	W S
Red Turtle Dove	<i>Streptopelia tranquebarica</i>	G	P	W
Racket Tailed Drongo	<i>Dicurus paradiseus</i>	G	I	W S
Red vented Bulbul	<i>Pycnonotus cafer</i>	G	O	W S
Red whiskered Bubul	<i>Pycnonotus jocosus</i>	G	O	W S
Red throated Flycatcher	<i>Ficedula parva</i>	G	I	W
Roseringed Parakeet	<i>Psittacula krameri</i>	G	P	W S
Rufous Treepie	<i>Dendrocitta vagabunda</i>	G	O	W S
Rufous Woodpecker	<i>Celeus brachyurus</i>	G	I	S
Scarlet Miiwet	<i>Pericrocotus flammeus</i>	W	I	W S
Shama	<i>Copsychus malabaricus</i>	W	I	W S
Small Minivet	<i>Pericrocotus cinnamomeus</i>	W	I	W S
Speckled Piculet	<i>Picumnus innominatus</i>	G	I	S

Spotted Dove	<i>Streptopelia chinensis</i>	W	P	W S
Streak throated woodpecker	<i>Picus xanthopygaeus</i>	Ev MD	I	S
Sulphur bellied Warbler	<i>Phylloscopus griseolus</i>	G	I	S
Tailor bird	<i>Orthotomus sutorius</i>	G	I	S
Thick billed Flowepecker	<i>Dicaeum agile</i>	Ev MD	P	W S
Tickell's Flycatcher	<i>Cyornis tickelliae</i>	W	P	W S
Tree Pippit	<i>Anthus trivialis</i>	G	I	S
Velvet fronted Nuthatch	<i>Sitta frontalis</i>	G	I	W S
Verditer Flycatcher	<i>Eumyias thalassinus</i>	Ev MD	I	W S
Vernal Hanging Parot	<i>Loriculus vernalis</i>	Ev MD	P	W S
Western Crown leaf Warbler	<i>Phylloscopus occipitalis</i>	Ev MD	I	W S
White bellied blue Flycatcher	<i>Cyornis pallipes</i>	Ev S	I	W S
White bellied Drongo	<i>Dicrurus caerulescens</i>	G	I	W S
White bellied Woodpecker	<i>Dryocopus javensis</i>	Ev MD	I	W S
White browed Bulbul	<i>Pycnonotus luteolus</i>	W	O	W
White browed Fantail	<i>Rhipidura aureola</i>	G	I	W
White checked Barbet	<i>Megalaima viridis</i>	G	P	W S
White rumped Munia	<i>Lonchura striata</i>	Ev MD	P	S
White throated Kingfisher	<i>Halcyon smyrnensis</i>	G	O	S
Yelllow Browed Bulbul	<i>Iole indica</i>	Ev MD	O	W S
Yellow Footed Green Pigeon	<i>Treron phoenicopterus</i>	G	P	W S

Habitat Guild

EvMD- Evergreen and Moist Deciduous Forest, W- Woodland, G- Generalist

Feeding Guild

I- Insectivore, P- Phytophagous, O- Other

Season

W- Winter, S- Summer

APPENDIX B- Birds seen in Sagar Forest Division

Common name	Scientific name
PODICIPEDIFORMES: Podicipedidae	
Little Grebe	<i>Tachybaptus ruficollis</i>
PELECANIFORMES: Phalacrocoracidae	
Indian Cormorant	<i>Phalacrocorax fuscicollis</i>
Great Cormorant	<i>Phalacrocorax carbo</i>
Little Cormorant	<i>Phalacrocorax niger</i>
PELECANIFORMES: Anhingidae	
Darter	<i>Anhinga melanogaster</i>
CICONIIFORMES: Ardeidae	
Great Egret	<i>Ardea alba</i>
Intermediate Egret	<i>Egretta intermedia</i>
Cattle Egret	<i>Bubulcus ibis</i>
Black-crowned Night-Heron	<i>Nycticorax nycticorax</i>
Gray Heron	<i>Ardea cinerea</i>
Purple Heron	<i>Ardea purpurea</i>
Little Egret	<i>Egretta garzetta</i>
Malayan Night-Heron	<i>Gorsachius melanolophus</i>
CICONIIFORMES: Ciconiidae	
Painted Stork	<i>Mycteria leucocephala</i>
Asian Openbill	<i>Anastomus oscitans</i>
Woolly-necked Stork	<i>Ciconia episcopus</i>
CICONIIFORMES: Threskiornithidae	
Black-headed Ibis	<i>Threskiornis melanocephalus</i>
Glossy Ibis	<i>Plegadis falcinellus</i>
Eurasian Spoonbill	<i>Platalea leucorodia</i>
ANSERIFORMES: Anatidae	
Lesser Whistling-Duck	<i>Dendrocygna javanica</i>
Ruddy Shelduck	<i>Tadorna ferruginea</i>
Comb Duck	<i>Sarkidiornis melanotos</i>
Cotton Pygmy-goose	<i>Nettapus coromandelianus</i>
Spot-billed Duck	<i>Anas acuta</i>
Northern Pintail	<i>Anas querquedula</i>
Garganey	<i>Anas clypeata</i>
Northern Shoveler	
FALCONIFORMES: Pandionidae	
Osprey	<i>Pandion haliaetus</i>
FALCONIFORMES: Accipitridae	
Oriental Honey-buzzard	<i>Pernis ptilorhynchus</i>
Black-shouldered Kite	<i>Elanus caeruleus</i>
Black Kite	<i>Milvus migrans</i>
Brahminy Kite	<i>Haliastur indus</i>

White-bellied Sea-Eagle	<i>Haliaeetus leucogaster</i>
Egyptian Vulture	<i>Neophron percnopterus</i>
White-rumped Vulture	<i>Gyps bengalensis</i>
Short-toed Eagle	<i>Circaetus gallicus</i>
Crested Serpent-Eagle	<i>Spilornis cheela</i>
Western Marsh-Harrier	<i>Circus aeruginosus</i>
Crested Goshawk	<i>Accipiter trivirgatus</i>
Shikra	<i>Accipiter badius</i>
Besra	<i>Accipiter virgatus</i>
White-eyed Buzzard	<i>Butastur teesa</i>
Tawny Eagle	<i>Aquila rapax</i>
Steppe Eagle	<i>Aquila nipalensis</i>
Bonelli's Eagle	<i>Aquila fasciata</i>
Booted Eagle	<i>Aquila pennata</i>
Changeable Hawk-Eagle	<i>Spizaetus cirrhatus</i>
FALCONIFORMES: Falconidae	
Eurasian Kestrel	<i>Falco tinnunculus</i>
Peregrine Falcon	<i>Falco peregrinus</i>
GALLIFORMES: Phasianidae	
Painted Francolin	<i>Francolinus pictus</i>
Gray Francolin	<i>Francolinus pondicerianus</i>
Red Spurfowl	<i>Galloperdix spadicea</i>
Gray Junglefowl	<i>Gallus sonneratii</i>
Indian Peafowl	<i>Pavo cristatus</i>
GRUIFORMES: Rallidae	
White-breasted Waterhen	<i>Amaurornis phoenicurus</i>
Ruddy-breasted Crake	<i>Porzana fusca</i>
Purple Swampfen	<i>Porphyrio porphyrio</i>
Common Moorhen	<i>Gallinula chloropus</i>
Eurasian Coot	<i>Fulica atra</i>
CHARADRIIFORMES: Jacanidae	
Pheasant-tailed Jacana	<i>Hydrophasianus chirurgus</i>
Bronze-winged Jacana	<i>Metopidius indicus</i>
CHARADRIIFORMES: Rostratulidae	
Greater Painted-snipe	<i>Rostratula benghalensis</i>
CHARADRIIFORMES: Recurvirostridae	
Black-winged Stilt	<i>Himantopus himantopus</i>
CHARADRIIFORMES: Glareolidae	
Small Pratincole	<i>Glareola lactea</i>
Red-wattled Lapwing	<i>Vanellus indicus</i>
Little Ringed Plover	<i>Charadrius dubius</i>
Lesser Sandplover	<i>Charadrius mongolus</i>
CHARADRIIFORMES: Scolopacidae	

Jack Snipe	<i>Lymnocyptes minimus</i>
Common Sandpiper	<i>Actitis hypoleucos</i>
Green Sandpiper	<i>Tringa ochropus</i>
Common Greenshank	<i>Tringa nebularia</i>
Marsh Sandpiper	<i>Tringa stagnatilis</i>
Wood Sandpiper	<i>Tringa glareola</i>
Common Redshank	<i>Tringa totanus</i>
Little Stint	<i>Calidris minuta</i>
CHARADRIIFORMES: Sternidae	
River Tern	<i>Sterna aurantia</i>
COLUMBIFORMES: Columbidae	
Rock Pigeon	<i>Columba livia</i>
Nilgiri Wood-Pigeon	<i>Columba elphinstonii</i>
Oriental Turtle-Dove	<i>Streptopelia orientalis</i>
Eurasian Collared-Dove	<i>Streptopelia decaocto</i>
Red Collared-Dove	<i>Streptopelia tranquebarica</i>
Spotted Dove	<i>Streptopelia chinensis</i>
Laughing Dove	<i>Streptopelia senegalensis</i>
Emerald Dove	<i>Chalcophaps indica</i>
Pompadour Green-Pigeon	<i>Treron pompadora</i>
Yellow-footed Pigeon	<i>Treron phoenicopterus</i>
Green Imperial-Pigeon	<i>Ducula aenea</i>
Mountain Imperial-Pigeon	<i>Ducula badia</i>
PSITTACIFORMES: Psittacidae	
Alexandrine Parakeet	<i>Psittacula eupatria</i>
Rose-ringed Parakeet	<i>Psittacula krameri</i>
Plum-headed Parakeet	<i>Psittacula cyanocephala</i>
Blossom-headed Parakeet	<i>Psittacula roseata</i>
Malabar Parakeet	<i>Psittacula columboides</i>
Vernal Hanging-Parrot	<i>Loriculus vernalis</i>
CUCULIFORMES: Cuculidae	
Common Hawk-Cuckoo	<i>Cuculus varius</i>
Indian Cuckoo	<i>Cuculus micropterus</i>
Banded Bay Cuckoo	<i>Cacomantis sonneratii</i>
Asian Koel	<i>Eudynamys scolopaceus</i>
Blue-faced Malkoha	<i>Phaenicophaeus viridirostris</i>
Sirkeer Malkoha	<i>Phaenicophaeus leschenaultii</i>
Greater Coucal	<i>Centropus sinensis</i>
STRIGIFORMES: Strigidae	
Collared Scops-Owl	<i>Otus lettia</i>
Rock Eagle-Owl	<i>Bubo bengalensis</i>
Spot-bellied Eagle-Owl	<i>Bubo nipalensis</i>
Brown Fish-Owl	<i>Ketupa zeylonensis</i>
Jungle Owlet	<i>Glaucidium radiatum</i>
Spotted Owlet	<i>Athene brama</i>
CAPRIMULGIFORMES: Podargidae	
Ceylon Frogmouth	<i>Batrachostomus moniliger</i>

CAPRIMULGIFORMES: Caprimulgidae	
Gray Nightjar	Caprimulgus indicus
Brown-backed Needletail	
Asian Palm-Swift	Hirundapus giganteus
Alpine Swift	Cypsiurus balasiensis
House Swift	Tachymarptis melba
Apus nipalensis	
APODIFORMES: Hemiprocnidae	
Crested Treeswift	Hemiproctne coronata
TROGONIFORMES: Trogonidae	
Malabar Trogon	Harpactes fasciatus
CORACIIFORMES: Alcedinidae	
Common Kingfisher	Alcedo atthis
Stork-billed Kingfisher	Pelargopsis capensis
White-throated Kingfisher	Halcyon smyrnensis
Black-capped Kingfisher	Halcyon pileata
Pied Kingfisher	Ceryle rudis
CORACIIFORMES: Meropidae	
Blue-bearded Bee-eater	Nyctyornis athertoni
Green Bee-eater	Merops orientalis
Blue-tailed Bee-eater	Merops philippinus
Chestnut-headed Bee-eater	Merops leschenaulti
CORACIIFORMES: Coraciidae	
Indian Roller	Coracias benghalensis
CORACIIFORMES: Upupidae	
Eurasian Hoopoe	Upupa epops
CORACIIFORMES: Bucerotidae	
Malabar Gray Hornbill	Ocyrceros griseus
Indian Gray Hornbill	Ocyrceros birostris
Malabar Pied-Hornbill	Anthracoceros coronatus
PICIFORMES: Capitonidae	
Brown-headed Barbet	Megalaima zeylanica
White-cheeked Barbet	Megalaima viridis
Crimson-fronted Barbet	Megalaima rubricapillus
Coppersmith Barbet	Megalaima haemacephala
PICIFORMES: Picidae	
Speckled Piculet	Picumnus innominatus
Brown-capped Woodpecker	Dendrocopos moluccensis
Yellow-crowned Woodpecker	Dendrocopos mahrattensis
Rufous Woodpecker	Celeus brachyurus
White-bellied Woodpecker	Dryocopus javensis
Lesser Yellownape	Picus chlorolophus
Streak-throated Woodpecker	Picus xanthopygaeus

Black-rumped Flameback	<i>Dinopium benghalense</i>
Greater Flameback	<i>Chrysocolaptes lucidus</i>
Heart-spotted Woodpecker	<i>Hemicircus canente</i>
PASSERIFORMES: Pittidae	
Indian Pitta	<i>Pitta brachyura</i>
PASSERIFORMES: Alaudidae	
Black-crowned Sparrow-Lark	<i>Eremopterix nigriceps</i>
Crested Lark	<i>Galerida cristata</i>
Malabar Lark	<i>Galerida malabarica</i>
Oriental Skylark	<i>Alauda gulgula</i>
PASSERIFORMES: Hirundinidae	
Dusky Crag-Martin	<i>Ptyonoprogne concolor</i>
PASSERIFORMES: Motacillidae	
Olive-backed Pipit	<i>Anthus hodgsoni</i>
Tree Pipit	<i>Anthus trivialis</i>
Forest Wagtail	<i>Dendronanthus indicus</i>
White-browed Wagtail	<i>Motacilla madaraspatensis</i>
Gray Wagtail	<i>Motacilla cinerea</i>
PASSERIFORMES: Campephagidae	
Large Cuckoo-shrike	<i>Coracina macei</i>
Black-headed Cuckoo-shrike	<i>Coracina melanoptera</i>
Small Minivet	<i>Pericrocotus cinnamomeus</i>
Scarlet Minivet	<i>Pericrocotus flammeus</i>
Bar-winged Flycatcher-shrike	<i>Hemipus picatus</i>
PASSERIFORMES: Pycnonotidae	
Black-crested Bulbul	<i>Pycnonotus melanicterus</i>
Red-whiskered Bulbul	<i>Pycnonotus jocosus</i>
Red-vented Bulbul	<i>Pycnonotus cafer</i>
White-browed Bulbul	<i>Pycnonotus luteolus</i>
Yellow-browed Bulbul	<i>Iole indica</i>
Black Bulbul	<i>Hypsipetes leucocephalus</i>
PASSERIFORMES: Chloropseidae	
Blue-winged Leafbird	<i>Chloropsis cochinchinensis</i>
Golden-fronted Leafbird	<i>Chloropsis aurifrons</i>
PASSERIFORMES: Aegithinidae	
Common Iora	<i>Aegithina tiphia</i>
PASSERIFORMES: Turdidae	
Blue-capped Rock-Thrush	<i>Monticola cinclorhynchus</i>
Blue Rock-Thrush	<i>Monticola solitarius</i>
Malabar Whistling-Thrush	<i>Myophonus horsfieldii</i>
Orange-headed Thrush	<i>Zoothera citrina</i>
Eurasian Blackbird	<i>Turdus merula</i>
White-bellied Shortwing	<i>Brachypteryx major</i>

PASSERIFORMES: Cisticolidae	
Zitting Cisticola	<i>Cisticola juncidis</i>
Jungle Prinia	<i>Prinia sylvatica</i>
Ashy Prinia	<i>Prinia socialis</i>
Plain Prinia	<i>Prinia inornata</i>
PASSERIFORMES: Sylviidae	
Blyth's Reed-Warbler	<i>Acrocephalus dumetorum</i>
Clamorous Reed-Warbler	<i>Acrocephalus stentoreus</i>
Booted Warbler	<i>Hippolais caligata</i>
Common Tailorbird	<i>Orthotomus sutorius</i>
Sulphur-bellied Warbler	<i>Phylloscopus griseolus</i>
Hume's Warbler	<i>Phylloscopus humei</i>
Greenish Warbler	<i>Phylloscopus trochiloides</i>
Western Crowned Leaf-Warbler	<i>Phylloscopus occipitalis</i>
PASSERIFORMES: Muscicapidae	
Asian Brown Flycatcher	<i>Muscicapa dauurica</i>
Red-breasted Flycatcher	<i>Ficedula parva</i>
Taiga Flycatcher	<i>Ficedula albicilla</i>
Verditer Flycatcher	<i>Eumyias thalassinus</i>
Nilgiri Flycatcher	<i>Eumyias albicaudatus</i>
White-bellied Blue-Flycatcher	<i>Cyornis pallipes</i>
Tickell's Blue-Flycatcher	<i>Cyornis tickelliae</i>
Gray-headed Canary-flycatcher	<i>Culicicapa ceylonensis</i>
Indian Blue Robin	<i>Luscinia brunnea</i>
Oriental Magpie-Robin	<i>Copsychus saularis</i>
White-rumped Shama	<i>Copsychus malabaricus</i>
PASSERIFORMES: Rhipiduridae	
White-browed Fantail	<i>Rhipidura aureola</i>
PASSERIFORMES: Monarchidae	
Black-naped Monarch	<i>Hypothymis azurea</i>
Asian Paradise-Flycatcher	<i>Terpsiphone paradisi</i>
PASSERIFORMES: Timaliidae	
Puff-throated Babbler	<i>Pellorneum ruficeps</i>
Indian Scimitar-Babbler	<i>Pomatorhinus horsfieldii</i>
Common Babbler	<i>Turdoides caudata</i>
Jungle Babbler	<i>Turdoides striata</i>
Yellow-billed Babbler	<i>Turdoides affinis</i>
Brown-cheeked Fulvetta	<i>Alcippe poiocephala</i>
Dark fronted Babbler	<i>Rhopocicla altriceps</i>
PASSERIFORMES: Paridae	
Great Tit	<i>Parus major</i>
Black-lored Tit	<i>Parus xanthogenys</i>
PASSERIFORMES: Sittidae	
Chestnut-bellied Nuthatch	<i>Sitta castanea</i>
Velvet-fronted Nuthatch	<i>Sitta frontalis</i>

PASSERIFORMES: Nectariniidae	
Purple-rumped Sunbird	<i>Leptocoma zeylonica</i>
Crimson-backed Sunbird	<i>Leptocoma minima</i>
Purple Sunbird	<i>Cinnyris asiaticus</i>
Western Crimson Sunbird	<i>Aethopyga vigorsii</i>
Little Spiderhunter	<i>Arachnothera longirostra</i>
Loten's Sunbird	<i>Cinnyris lotenius</i>
PASSERIFORMES: Dicaeidae	
Thick-billed Flowerpecker	<i>Dicaeum agile</i>
Pale-billed Flowerpecker	<i>Dicaeum erythrorhynchos</i>
Plain Flowerpecker	<i>Dicaeum concolor</i>
PASSERIFORMES: Zosteropidae	
Oriental White-eye	<i>Zosterops palpebrosus</i>
PASSERIFORMES: Oriolidae	
Eurasian Golden Oriole	<i>Oriolus oriolus</i>
Black-naped Oriole	<i>Oriolus chinensis</i>
Black-hooded Oriole	<i>Oriolus xanthornus</i>
PASSERIFORMES: Irenidae	
Asian Fairy-bluebird	<i>Irena puella</i>
PASSERIFORMES: Laniidae	
Brown Shrike	<i>Lanius cristatus</i>
Long-tailed Shrike	<i>Lanius schach</i>
PASSERIFORMES: Prionopidae	
Large Woodshrike	<i>Tephrodornis gularis</i>
Common Woodshrike	<i>Tephrodornis pondicerianus</i>
PASSERIFORMES: Dicuridae	
Black Drongo	<i>Dicurus macrocerus</i>
Ashy Drongo	<i>Dicurus leucophaeus</i>
White-bellied Drongo	<i>Dicurus caerulescens</i>
Bronzed Drongo	<i>Dicurus aeneus</i>
Hair-crested Drongo	<i>Dicurus hottentottus</i>
Greater Racket-tailed Drongo	<i>Dicurus paradiseus</i>
PASSERIFORMES: Artamidae	
Ashy Woodswallow	<i>Artamus fuscus</i>
PASSERIFORMES: Corvidae	
Rufous Treepie	<i>Dendrocitta vagabunda</i>
White-bellied Treepie	<i>Dendrocitta leucogastra</i>
House Crow	<i>Corvus splendens</i>
Large-billed Crow	<i>Corvus macrorhynchos</i>
PASSERIFORMES: Sturnidae	
Common Hill Myna	<i>Gracula religiosa</i>
Jungle Myna	<i>Acridotheres fuscus</i>

Common Myna	<i>Acridotheres tristis</i>
Asian Pied Starling	<i>Gracupica contra</i>
Chestnut tailed starling	<i>Sturnia malabarica</i>
PASSERIFORMES: Passeridae	
House Sparrow	<i>Passer domesticus</i>
Chestnut-shouldered Petronia	<i>Petronia xanthocollis</i>
PASSERIFORMES: Estrildidae	
White-rumped Munia	<i>Lonchura striata</i>
PASSERIFORMES: Fringillidae	
Common Rosefinch	<i>Carpodacus erythrinus</i>